

ISSN 0147-9725



MARYLAND BIRDLIFE

Avian studies in the Mid-Atlantic region



Bulletin of the Maryland Ornithological Society, Inc.

**SPRING 2017
VOLUME 66
NUMBER 1**

MARYLAND ORNITHOLOGICAL SOCIETY, INC.

P.O. Box 105, Monrovia, MD 21770-0105

301-588-4250

<http://www.mdbirds.org>

CURRENT OFFICERS – SPRING 2017

EXECUTIVE COUNCIL

| | | | |
|-----------------|-----------------|--|--------------|
| President: | Barbara Johnson | barbarajohnson222@gmail.com | 410-703-4664 |
| Vice-President: | Robin Todd | totnesman@aol.com | 410-491-5333 |
| Treasurer: | Dave Webb | porzana@comcast.net | 410-939-3537 |
| Secretary: | Maryanne Dolan | maryanne.dolan@gmail.com | 443-350-9197 |
| Past-President: | Tom Strikwerda | tstrik@verizon.net | 301-942-2841 |

STATE DIRECTORS

| | | | |
|---------------------|---|-------------------|---|
| Allegany & Garrett: | *Mary Huebner (vacant) | Howard: | *John Harris Karen Heffernan John McKitterick Robin Todd |
| Anne Arundel: | *David Gillum Diane Benyus Barbara Ricciardi | Kent: | *Nancy Martin Walter Ellison |
| Baltimore: | *Peter Lev Joan Cwi Mary Anne Fluke Mark Linardi | Montgomery: | *Clive Harris Andy Martin Jim Moore Brooke Smith |
| Caroline: | *Danny Poet (vacant) | Patuxent: | *Fred Fallon Marcia Watson |
| Carroll: | *Don Jewell (vacant) | Talbot: | *Dave Palmer Wayne Bell |
| Cecil: | *Maryanne Dolan John Taylor | Tri-County: | *Mike Walsh Rick Palmer |
| Frederick: | *Kathy Brown Tom Humphrey | Washington (Co.): | *George Warrick Mark Abdy |
| Harford: | *Bob Werrlein Thomas Congersky Dennis Kirkwood | | |

**Denotes Chapter President*

Active Member: \$20.00 plus chapter dues

Household: \$25.00 plus chapter dues

Sustaining: \$50.00 plus chapter dues

Life: \$1000.00 (4 annual installments)

Junior (under 18): \$5.00 plus chapter dues

Cover: Long-eared Owl, *Asio otus*. Delmarva Peninsula, Worcester County, Maryland, 8 November 2008. Photographed by George M. Jett©.

EDITOR'S NOTE

Chandler S. Robbins: The Passing of an Ornithological Icon

It is with great sadness that I report the passing of Chandler S. Robbins on 20 March 2017 at the age of 98. Chan edited *Maryland Birdlife* from 1947 through 2014. A chronicle of his amazing life can be found in the spring 2015 issue of *Maryland Birdlife*. The current issue is dedicated to Chan's memory.

Latest AOS Checklist of North and Middle American Birds

This issue of *Maryland Birdlife* will utilize the American Ornithological Society (AOS; formerly AOU) Checklist of North and Middle American Birds through the Fifty-seventh Supplement (2016). The list order has changed extensively. Details of the changes can be found in:

Chesser, R.T., K.J. Burns, C. Cicero, J.L. Dunn, A.W. Kratter, I.J. Lovette, P.C. Rasmussen, J.V. Remsen, Jr., J.D. Rising, D.F. Stotz, and K. Winker. 2016. Fifty-seventh Supplement to the American Ornithologists' Union Check-list of North American Birds. *The Auk: Ornithological Advances* 133:544–560. Available at:
<http://americanornithologypubs.org/doi/pdf/10.1642/AUK-16-77.1>.

New Statewide May Count Compiler

It is my pleasure to introduce Marilyn Veek as the Maryland Ornithological Society's new statewide May Count Compiler. The following is in Marilyn's words:

"There is nothing like the thrill of getting a great look at a new bird which previously only existed as a plate in a bird guide. On the other hand, birding the same areas repeatedly over seasons and years, whether my own property, a nearby park, or a sector on an annual count, is its own kind of pleasure—seeing what is new, what is the same, whether the White-throated Sparrows have appeared as a beacon of winter's approach. I particularly enjoy birding travel to new parts of the world; learning a bit about a country's culture and geography while watching its beautiful birds is a wonderful experience. I've been a member of the Frederick Bird Club for over 20 years, and currently serve as the Secretary. I am also the Calendar Editor for *The Maryland Yellowthroat*, which has brought me into contact with members from MOS chapters all over the state, as well as has sharpened my editing skills! I learned so much about birds and birding from Mike Welch during club meetings and field trips, that volunteering to serve as May Count Compiler when he moved to Arizona seemed only appropriate."

Eugene J. Scarpulla
Editor
ejscarp@comcast.net

Long-eared Owl (*Asio otus*) Annual Frequency, Seasonal Chronologies, and Status on the Delmarva Peninsula

Jan G. Reese^{1,3} and Robbie Weterings²

¹*P.O. Box 298, St. Michaels, Maryland 21663*

²*Forests and Climate Change Programme, German Society for International Collaboration (GIZ), Jakarta, Indonesia*

³*Corresponding author: reesejan@ymail.com*

Abstract: We determined the occurrence distribution, annual frequency, and seasonal chronologies of Long-eared Owls (*Asio otus*) on the Delmarva Peninsula using various types of records from 1941 through 2016. Records were distributed throughout the peninsula with annual peak frequencies fluctuating in a general 3–5-year periodicity. Seasonal records occurred in three periods where the frequency increased and subsequently decreased. An early-March to mid-April period approximating northbound migrants peaked on 19 March (± 3.2 SEM), a late-October to mid-December period approximating southbound migrants peaked on 14 November (± 2.4 SEM), and a late-November to late-March winter period peaked on 12 January (± 2.6 SEM). Historically there is no record of nesting on the peninsula, nor was it confirmed among the few spring and summer records in this study. The Long-eared Owl is a poorly understood species with little to nothing known about it regionally. This study provides some valuable insights to their life history on the Delmarva Peninsula, a more definitive status than that given for the region in the literature, and furnishes baseline information for future study.

Keywords: annual frequency, *Asio otus*, Delmarva Peninsula, distribution, Long-eared Owl, seasonal chronologies, status

Long-eared Owl (*Asio otus*) occurrence and nesting distribution east of the Great Lakes in North America is associated with the contiguous mountain regions north from central Pennsylvania through the Canadian Maritime Provinces. Records of the species occurring and nesting farther east and south are considered rare and local (Marks et al. 1994). Most owls withdraw in winter from northern parts of their range and may be found occasionally wintering as far south as central Georgia and southern Mississippi (Marks et al. 1994, Beaton et al. 2003, Mississippi Ornithological Society 2015).

Long-eared Owls are infrequently detected owing, to their secretive behavior with faint calls, their more post-dusk nocturnal activity in comparison to other

owl species, their superb cryptic plumage coloration and posturing when perched, and temporary seasonal occurrence in southeast regions with abundant prey and lack of prolonged snow cover. These factors make the species a difficult study subject and therefore data concerning their behavior, biology and phenology are limited. This may obscure the actual status of the species throughout their eastern North America range including the Delmarva Peninsula where the species is considered scarce/rare.

Additional factors complicate determining the actual status of Long-eared Owls in the region. One factor is the “broad-scale” at which the status for a species is sometimes assigned. Portions of three political jurisdictions (Delaware, Maryland, and Virginia) comprise the Delmarva Peninsula Coastal Plain. The jurisdictions contain various other physiographic regions to the north, west, and south. Most studies attribute a given species status to the entire political jurisdiction, and not to the individual physiographic regions within (Rives 1890, Kirkwood 1895, Pennock 1904, Rhoads and Pennock 1905, Hampe and Kolb 1947, Murray 1952). Few if any species are found equally dispersed across all physiographic regions due to variation in habitat and food availability within the different regions. Thus, assigning a species status to an entire geographic area, when that area’s boundaries are not based on any physiographic or ecological context, may be misleading and create an inaccurate picture of the owl’s status in the jurisdictional region.

Historical studies from around the turn of the 20th Century considered Long-eared Owls to be “uncommon” or “resident” on the Delmarva Peninsula despite more abundant concealing habitat and fewer observers to detect owls relative to more modern times (Rives 1890, Kirkwood 1895, Pennock 1904, Rhoads and Pennock 1905). Conversely, studies during the mid-20th Century classified the species status as “rare” (Hampe and Kolb 1947, Murray 1952, Stewart and Robbins 1958). Later studies associated the perceived less frequent detection and/or population decline to a mid-century surge in Great Horned Owl (*Bubo virginianus*) and Red-tailed Hawk (*Buteo jamaicensis*) populations (Bosakowski et al. 1989, Hess et al. 2000). These two species were afforded legal protection after a century of persecution by farmers and hunters. Competition for nest sites and small rodent prey by the Great Horned Owl and other raptors has been suggested as possibly limiting Long-eared Owl numbers in open lands of New Jersey (Bosakowski et al. 1989). There were also reports of Great Horned Owls and Red-tailed Hawks preying/attacking Long-eared Owls (Collins 1962, Bosakowski et al. 1989). It is uncertain whether these factors may have contributed to mixed status reports of “rare to uncommon” for studies including the Delmarva Peninsula in the 21st Century (Hess et al. 2000, Rottenborn and Brinkley 2007).

In this paper, we used regional publications, checklists, reports, seasonal counts, banding records and museum specimens of Long-eared Owl occurrences on the Delmarva Peninsula since the early 1900s to compile a comprehensive dataset. These data were then used to determine annual frequency, seasonal chronologies and status on the peninsula.

METHODS

Study Area

The Delmarva Peninsula Coastal Plain is located in the Mid-Atlantic region of North America and comprises a peninsula situated between the Atlantic Ocean on the east and Chesapeake Bay on the west (Figure 1). The peninsula is approximately 290 km (180 mi) long with a north/south aspect, nearly 110 km (68 mi) wide midway its length and tapers towards both ends. Alluvial deposits of silt, sand, clay, and gravel from glacier melt during the Miocene, Pleistocene, and Holocene epochs form the peninsula's substrates. The topography lacks pronounced relief and elevation is generally less than 30 m (98 ft) above sea level. Freshwater streams and rivers, brackish and saltwater tidal creeks, tributaries and bays deeply indent the landscape of prevalent cultivated agricultural fields while coniferous and deciduous forests cover most of the undeveloped areas. Other areas are characterized by herbaceous and forested freshwater wetlands, and herbaceous brackish and saltwater wetlands. Additionally linear barrier islands and bays characterize the coast with the Atlantic Ocean. Average temperatures range is -2.8° to 7.8° C (27° to 46° F) in January and 18.3° to 30.5° C (65° to 87° F) in July at the center of the peninsula. Annual precipitation averages 1143 mm (45 in), and snowfall, 533 mm (21 in). The humidity ranges 54% to 78% while prevailing west winds average 16.1 to 17.7 km/hr (10 to 11 mi/hr). The study area encompasses the entire Coastal Plain peninsula south of its interface with the Piedmont upland physiographic region.

Data Sources

Long-eared Owl records were extracted from sources given in Table 1, plus regional bird books and checklists cited herein. Of these records, two were found to be duplicates within the dataset, one gave no location, one was only reported in an unreliable source, and one museum inventory specimen could not be found in the collection. These records date from the last half of the 20th Century, could not be validated, and were therefore deleted from the dataset. All remaining records used in the dataset occurred during the period 1941–2016. Additionally, private individuals (noted in the acknowledgments) contributed Long-eared Owl records from their personal observations, counts, and/or records.




Figure 1. Delmarva Peninsula study area showing the distribution of 50 locations where 218 Long-eared Owls were recorded from 1941 through 2016 (some locations are represented by more than one dot).

Table 1. Publications, museum specimen inventories, and databases used in compilation of the Long-eared Owl dataset.

| Journals and Databases | Data Period | URL |
|--|-----------------------|--|
| <i>Cassinia</i> | 1890–2009 | http://dvoc.org/Publications/Cassinia/Cassinia.htm |
| <i>Delmarva Ornithologist</i> | 1964–1974 | http://olddos.dosbirds.org/ornithologists |
| <i>Maryland Birdlife</i> | 1945–2016 | https://sora.unm.edu/node/132691 (2006–2016 hardcopy subscription) |
| <i>The Raven</i> | 1930–2013 | http://www.ccbbirds.org/resources/the-raven/ |
| <i>North American Birds</i> | 1999–2008 | https://sora.unm.edu/node/209 |
| <i>National Audubon Society Field Notes</i> | 1994–1998 | https://sora.unm.edu/node/209 |
| <i>American Birds</i> | 1973–1994 | https://sora.unm.edu/node/209 |
| National Audubon Society Christmas Bird Counts | 1941–2014 | http://netapp.audubon.org/cbcobservation/ |
| eBird.org (Explore Data) | Through 1 Mar 2016 | http://ebird.org/ebird/explore |
| <i>List of Caroline County Birds</i> | 1946–1978 | Fletcher et al. 1956, Fletcher et al. 1979 |
| USGS Bird Banding Laboratory | 1941–2016 | Unpublished data retrieved 15 SEP 2016, Patuxent Wildlife Research Center, Laurel, MD, USA |

| Museum and University | Bird Collection Inventories | Access Date | Data Sources |
|---|------------------------------------|--------------------|---|
| Yale University | | 1 Jan 2016 | http://collections.peabody.yale.edu/search/ |
| American Museum of Natural History | | 1 Jan 2016 | http://sci-web-001.amnh.org/db/emuwebamnh/index.php |
| Philadelphia Academy of Science | | 5 Jan 2016 | http://www.ansp.org/research/systematics-evolution/collections/ornithology/ |
| Delaware Museum of Natural History | | 5 Jan 2016 | http://www.vertnet.org/ |
| Towson University | | | personal 1988 hard copy inventory (and L. Scott Johnson, in litt., 2010) |
| Salisbury University | | | personal 2015 hard copy inventory |
| University of Maryland (Horn Point & College Park campuses) | | | personal 1998 hard copy inventories |
| U.S. National Museum of Natural History | | 8 Jan 2016 | http://collections.nmnh.si.edu/search/birds/ |

Data Analysis

We converted all records to presence-only data to account for methodological differences among records from the various data sources. These records consisted of single or multiple owls encountered at a single location. In five cases, single or multiple owls were seen at a single location on widely separate dates (>1 weeks). For these observations the first and the last dates were included, e.g., the first and last dates seen at a winter roost site. The locations of all records were geocoded and visualized on a map using ArcGIS 10.1 (Environmental Systems Research Institute 2012). We calculated the total number of records per year for the period 1941–2016 and plotted them to display the annual frequency. We grouped all records from different years to calculate the number of records per week and plotted them to visualize seasonal variation throughout the year. We excluded Christmas Bird Counts from the latter analysis because their inclusion would have disproportionately skewed wintering records falsely high during the annual count period. Using the plotted seasonal records, we identified three clusters during which the number of records increased and subsequently decreased (approximating mid-February to mid-April, late-October to early-December, and mid-December to mid-February), plus six scattered records during May–September. These periods suggest northbound and southbound migration and wintering periods with each period partly overlapping the adjoining period.

To better understand the extent of these periods and overlaps, we assigned each record to one of the seasonal periods. Records of single or multiple owls at historical or known roosts and all owl records in January to mid-February (wintering owls) were assigned to a late-November to late-March period; records of single owls found at random non-roost locations were assigned to either the northbound early-March to mid-April period or the southbound late-October to mid-December period depending on whether they occurred before or after the peak wintering months of January to mid-February. To assure compliance and accuracy we compared the range of these periods with published early/late arrival dates for migrant and wintering owls given for jurisdictions in the Mid-Atlantic region. Only six records were available for the May–September period, hence this period was excluded from analysis.

Finally, we calculated the mean date during which the number of records was highest for the three different periods. Dates were first converted to Julian days in order to ease calculations. The mean dates were then plotted together at the 95% confidence interval (2.5 and 97.5 percentiles). All analyses were conducted in RStudio Version 0.99.491 (RStudio 2015) built on R 3.2.0 (R Development Core Team 2016); plots were created using the ggplot package.

RESULTS

Composition of Records

We extracted and considered 218 valid Long-eared Owl records from the Delmarva Peninsula in 1941–2016. The annual Christmas Bird Counts comprised 121 owl records during the count period 14 December through 5 January involving 18 different count locations. An additional 97 records from 32 locations included 61 sightings other than Christmas Bird Count dates, 12 trapped and banded, eight found dead on the highway, eight heard calling, three purposely shot for collections, three found injured, one found sick, and one tangled in a fence.

Distribution of Records

The early-March to mid-April period records ($n = 15$) were distributed across the northern portion of the peninsula north of a line from Cape Henlopen at the mouth of Delaware Bay west to the mouth of the Choptank River on Chesapeake Bay with exception of one record on the north side of the mouth of the Pocomoke River on Chesapeake Bay (Figure 1). Conversely, 85% of the late-October to mid-December period records ($n = 26$) were evenly dispersed at locations south of a line from Cape Henlopen on Delaware Bay west to Eastern Bay on Chesapeake Bay. The late-November to late-March period records ($n = 171$ including 121 Christmas Bird Counts) were broadly distributed with 81% adjacent to Delaware Bay ($n = 60$), Chesapeake Bay ($n = 33$), and Atlantic Ocean ($n = 45$) tidewater, and the remaining at more inland locations ($n = 33$). There were six May–September period records all distributed north of a line from Cape Henlopen at the mouth of Delaware Bay west to the mouth of the Choptank River on Chesapeake Bay.

Annual Records

Long-eared Owls have been reported on the Delmarva Peninsula every year since 1964 (52 years) with the exception of 1972 and 1985 (Figure 2). Annual records since the mid-1940s fluctuated generally reaching highs every 3–5 years before an appreciable drop in the number of records the following year(s).




Figure 2. Long-eared Owl annual frequency of records ($n = 218$) on the Delmarva Peninsula.

Seasonal Records

Figure 3 shows the Long-eared Owl seasonal records on the Delmarva Peninsula. The early-March to mid-April period records ($n = 15$) ranged from the 2 March early arrival through the 9 April late arrival with a mean peak date for the number of records on 19 March (± 3.2 SEM) (Figure 3B). The late-October to mid-December period records ($n = 26$) ranged from the 23 October early arrival through the 17 December late arrival with a mean date on 14 November (± 2.4 SEM). The late-November to late-March period records ($n = 50$), excluding Christmas Bird Counts, spanned 24 November through 24 March with a mean peak date on 12 January (± 2.6 SEM).




Figure 3. Long-eared Owl seasonal records on the Delmarva Peninsula.

A) Each vertical bar represents the number of records found in a week using data for the period 1941–2016 excluding Christmas Bird Counts. B) The vertical line in the box is the mean peak date for the period, the gray represents the confidence interval limits and line whiskers are the period extremes.

Age and Sex of Owls

Age and/or sex given in 28 records of Long-eared Owls trapped and banded, found sick or dead, or purposely collected, designated 16 (57%) as age unknown. Females comprised 85% of 13 owls for which the sex was determined and females were predominant among the migratory and wintering period owls. The age and sex for each of the four seasonal periods are summarized in Table 2.

Table 2. Long-eared Owl age and sex taken from seasonal records on the Delmarva Peninsula during 1941–2016. M = male, F = female, U = unknown sex.

| | early-Mar–mid-Apr period (n = 9) | | | late-Oct–mid-Dec period (n = 10) | | | late-Nov–mid-Mar period (n = 7) | | | May–Sep period (n = 2) | | |
|-------------|-------------------------------------|---|---|-------------------------------------|---|---|------------------------------------|---|---|---------------------------|---|---|
| Sex | M | F | U | M | F | U | M | F | U | M | F | U |
| Age <1 | - | - | - | - | 1 | 2 | - | - | - | - | - | - |
| Age >1 | - | 2 | - | - | 4 | 1 | 1 | - | 1 | - | - | - |
| Age unknown | - | 1 | 6 | - | - | 2 | - | 2 | 3 | 1 | 1 | - |

May–September Records

Owls calling during the first half of May at locations adjacent to the west shore of Delaware Bay in the northeast portion of the peninsula constituted three May–September records, while others included single owls found sick on a lawn in June and dead on the highway in July adjacent to the east side of Chesapeake Bay in the west-central portion of the peninsula, and another single reported occurring in September near the center of the peninsula.

DISCUSSION

Despite their infrequent detection and perceived rare status, 1941–2016 records of Long-eared Owls on the Delmarva Peninsula indicate they occur throughout the peninsula during migration and/or winter periods more frequently than generally thought. Additionally, there are a few warm-weather records, but no nest confirmations.

Distribution of Records

Most early-March to mid-April period records were distributed in the north portion of the peninsula while the late-October to mid-December period records favored the central and southern portions. Hawk watchers, banding stations, and bird watchers focus along the Atlantic and Chesapeake coasts during the fall migration season, which may have contributed to the seasonal difference found in the distribution. Owls found on the annual Christmas Bird Counts during the study contributed substantially to the late-November to late-March period records distribution.

Annual Records

Periodic spikes in annual frequency (generally every 3–5 years) noted in this study were also found in Long-eared Owl reports in New Jersey during 1966–1986 (Bosakowski et al. 1989). Periodic spikes were also found among banding station records in New York during 1981–1986 and New Jersey during 1980–

1988 (Slack et al. 1987, Duffy and Kerlinger 1992). In the northern boreal forest of eastern North America, and in northern Scotland and Fennoscandia such changes in owl frequency have been associated with low abundance of voles and lemmings in their usual distribution range, e.g., meadow voles, *Microtus* spp. (Harvey and Riddiford 1990); *Microtus* spp. and lemmings, *Lemmus* spp. (Korpimäki and Krebs 1996); and the Southern Red-backed Vole, *Myodes gapperi* (Cheveau et al. 2004). Fluctuations in Long-eared Owl abundance and reproductive success have been positively correlated to density of its primary microtine prey in Scotland (Field Vole, *Microtus agrestis* [Village 1981]) and western Finland (*M. agrestis* and Southern Vole, *M. levis* [Korpimäki and Norrdahl 1991]). Similarly, the closely related Short-eared Owl (*Asio flammeus*) exhibited a positive correlation between the abundance of voles/lemmings and owls in Alaska (Brown Lemming, *L. sibiricus* [Pitelka et al. 1955a, 1955b]), Scotland (*M. agrestis* [Village 1987]), and Finland (*M. agrestis* and *M. levis* [Korpimäki and Norrdahl 1991]). Monitoring prey was beyond the scope of this study, therefore we cannot with any certainty assign the periodicity to prey availability. Nevertheless, prey availability is a likely factor in driving the patterns in our data as is seen for Long-eared Owl in other studies. Finally, decreasing owl habitat with increasing observers utilizing improved detection technologies may have been responsible for the generally increasing annual frequency of owl records after the mid-20th Century.

Seasonal Records

Christmas Bird Counts frequently involve concerted efforts to locate rare species such as Long-eared Owls in the days just prior to the count and confirming their presence on count day. Therefore, Christmas Bird Counts were excluded from the seasonal analysis to avoid biasing the records peak for the wintering period during the 14 December through 5 January Christmas Bird Count period. The three periods seen in the data suggest migrant and wintering periods. The early-March to mid-April period records in this study correspond to northbound migrant Long-eared Owl capture periods of 29 March to 26 April and 21 March to 14 April at banding stations along the west and east portions of the south shore of Lake Ontario in New York (Beardslee and Mitchell 1965, Slack et al. 1987). The late-October to mid-December period records correspond with the 11 October to 27 November period of banding captures at the nearby Cape May Point, New Jersey banding station in 1980–1989 (Duffy and Kerlinger 1992). Owls banded at this station were rarely captured twice in a season suggesting they were south migrants and not resident birds (Katherine E. Duffy, in litt.). Also, a nocturnal visual study of southbound migrant owls at Cape May in 1982 detected owls most frequently during 12 October to 11 November (Russell et al. 1991). The late-October to mid-December period records also correspond to dates (10 October–18 November) for early southbound migrants on the Virginia Coastal Plain (Larner 1979, Kain 1987, Rottenborn and Brinkley 2007). All of

these seasonal chronologies lend support to Long-eared Owls being an annual migrant on the Delmarva Peninsula.

Despite evidence supporting the three periods as representing migrant and wintering periods, it is not without some caution. The seasonal analysis indicates an early-March to late-March overlap of 23 days between the late-November to late-March period records and the early-March to mid-April period records. Similarly there is a late-November to mid-December overlap of 24 days between the late-October to mid-December period records and the late-November to late-March period records. It is difficult to ascertain owl records to migrants or wintering. Furthermore, nest activity has been reported from mid-March into July in adjacent Piedmont and Ridge and Valley physiographic regions (Santner 1992, Rottenborn and Brinkley 2007, Ellison 2010, Gross 2012), while nothing is known about regional owl movements in winter. Regardless, this study contributes to further understanding of the migration and winter chronologies of this poorly understood species.

May-September Records

Historically, the Long-eared Owl was listed as “resident” in Delaware (Rhoads and Pennock 1905) and Virginia (Rives 1890), and “resident, but...not common” and “rare permanent resident” in Maryland (Kirkwood 1895, Hampe and Kolb 1947) which implies potential nesting on the Delmarva Peninsula. Regional studies since the first half of the 20th Century (Murray 1952, Stewart and Robbins 1958, Hess et al. 2000, Rottenborn and Brinkley 2007, Ellison 2010) restrict any potential nesting in those jurisdictions to Piedmont or higher elevations north, south, or west of the Coastal Plain peninsula. Nesting was documented just 3 km (2 mi) east of the Delmarva Peninsula on the New Jersey Coastal Plain near Salem (Stone 1965); 13 km (8 mi) north on the Delaware Piedmont north of Wilmington (B.C. Hiatt in Potter 1937), and 52 km (32 mi) west on the Maryland Piedmont near Baltimore (Kirkwood 1895, Kolb 1947a, Kolb 1947b, Stewart and Robbins 1958). Coincidentally, the three 21st Century owl calling records in May are on the west side of Delaware Bay just 12–32 km (7.5–20 mi) southeast of the late 1930s Coastal Plain nesting near Salem, New Jersey and the Piedmont nesting north of Wilmington, Delaware.

Necropsy of a male owl, found sick on a lawn, in June 1981 (JGR catalogue no.1045, USNM cat. No. 582672), and a female, found dead on the highway, in July 1983 (JGR catalogue no. 1400, USNM cat. No.597098) lacked frayed feathers, enlarged gonads, incubation patches, or other signs of nesting. It should be noted however, that with mid-March initiation of nest activity (Stewart and Robbins 1958, Marks et al. 1994, Ellison 2010) and subsequent egg-laying, incubation, and brooding of nestlings could have been completed by May and thereafter exhibit no physical signs of nesting. An owl reported on 14 September 1941 near the center of the peninsula suggests a resident or unusually early

southbound migrant. Clark (1975) found that Short-eared Owls shift their wintering and breeding areas in response to spatial and temporal abundance of small rodents. Perhaps small mammal population dynamics contributed to the occurrence of six spring and summer Long-eared Owl records. Indeed, all of the records coincide with years of high owl abundance on the peninsula. In summary, Long-eared Owl nesting has not been confirmed on the Delmarva Peninsula since the turn of the 20th Century despite several spring and summer records.

Owl Age and Sex

Handling of 28 Long-eared Owls trapped and banded, found sick or dead, or purposely collected enabled the possibility of determining the age and/or sex of individuals. Age determined for 12 November–April owls included nine greater than one-year-old (75%) and three less than one-year-old. Similarly, an overwhelming 97% of 37 northbound migrant Long-eared Owls banded at Nine Mile Point, New York in 1981–1986 (Slack et al. 1987) were greater than one-year-old. Conversely, only 26% of 203 southbound migrants banded at Cape May Point in 1980–1988 (Duffy and Kerlinger 1992) were greater than one-year-old. Age was not determined for any May–September owls. It should be noted that reliable external criteria for determining Long-eared Owl age/sex were lacking or limited prior to publication of the *Identification Guide to North American Birds* (Pyle 1997) that cautions age determination may be unreliable for some Long-eared Owls less than one-year-old or in the following year.

Sex determined for 13 owls in this study included 11 females (85%). In Britain, females constituted 78% of 36 Long-eared Owl southbound migrants banded at Fair Isle, Shetland, Scotland in 1984–1989 (Harvey and Riddiford 1990) while females comprised 76% of 101 owl mortalities autopsied in the United Kingdom during October–April 1963–1995 (Wyllie et al. 1996). Conversely, females comprised only 36% of 36 winter mortalities in Norway (Overskaug and Kristiansen 1994). In Fennoscandia, males winter close to their nest site and territory to assure retaining them for the following spring while females have no constraints and move to other areas in times of food shortage (Harvey and Riddiford 1990). It is not known if this type of movement contributed to the over-representation of females in this study while this domain remains largely unexplored for the Long-eared Owl. In addition, the sex ratios in our study and the southbound migrants banded in Scotland were based on small sample sizes while the contradictory ratio came from a Norway study 2414 km (1500 mi) north of this study. Therefore, it is difficult to make any conclusions about age and sex ratios in our study.

Status

The species general status given in late 19th Century into early 21st Century literature ranges from “rare resident” in the north portion of the peninsula to “rare resident, transient or wintering” in the southern portion (Table 3).

Table 3. Long-eared Owl historical status, occurrence and nesting on the Delmarva Peninsula Coastal Plain.

| Political Jurisdiction | Status | Occurrence | Nesting | Source |
|------------------------|--|---|---|------------------------------|
| Delaware | Throughout the year (statewide) | More abundant during cold months | Occasionally breeds within our limits | Pennock 1904 |
| Delaware | Resident (statewide) | N/A | N/A | Rhoads and Pennock 1905 |
| Delaware | Regular winter visitor, uncommon (statewide) | Nov-Mar | Newcastle County 1937; today DE should be considered south of normal breeding range | Hess et al. 2000 |
| Maryland | Resident, not common (statewide) | No records for Coastal Plain | No records for Coastal Plain | Kirkwood 1895 |
| Maryland | Rare permanent resident (statewide) | No records for Coastal Plain | No records for Coastal Plain | Hampe and Kolb 1947 |
| Maryland | Rare and local permanent resident west of Chesapeake Bay, occasionally occurs on Coastal Plain | Fall and winter in Dorchester and Caroline Counties | No records for Coastal Plain | Stewart and Robbins 1958 |
| Maryland | Year-round resident in nesting range (statewide) | N/A | No records for Coastal Plain | Ellison 2010 |
| Virginia | Resident, but seldom seen (statewide) | Occurrence not defined | No records for Coastal Plain | Rives 1890 |
| Virginia | Rare resident (statewide) | Recorded in winter most often | No records for Coastal Plain | Murray 1952 |
| Virginia | Rare winter visitor (Coastal Plain) | 18 Nov–20 May | No records for Coastal Plain | Larner 1979 |
| Virginia | Rare winter visitor (Coastal Plain) | 31 Oct–28 May | No records for Coastal Plain | Kain 1987 |
| Virginia | Rare transient and winter visitor (Coastal Plain) | 10 Oct–28 May | No records for Coastal Plain | Rottenborn and Brinkley 2007 |

Recent Wildlife Action Plans for jurisdictions comprising the Delmarva Peninsula rank the Long-eared Owl status as “highly state rare breeding and non-breeding” (S1B, S1N) in Delaware (Delaware Division of Fish and Wildlife 2015) and in Maryland (Maryland Department of Natural Resources 2015). In addition Delaware provides an SHB rank, i.e., known historically to breed in Delaware. Virginia provides no ranking for the species on either the Accomack or Northampton County Coastal Plain (Virginia Department of Game and Inland

Fisheries 2015). In view of the variable statuses applied to inclusive multi-physiographic jurisdictions during the past 74 years, it is no wonder status of the species is clouded and/or contradictory. This study found records of Long-eared Owls on the Delmarva Peninsula show regular annual frequency with seasonal chronologies during migrant and wintering periods while fluctuating abundance may decrease the number of records in some years.

Conclusion

This study of 1941–2016 Long-eared Owl records on the Delmarva Peninsula found them distributed throughout the peninsula with annual peak frequencies fluctuating in a general 3- to 5-year periodicity. Seasonal records show three distinct periods approximating migrant and wintering periods. Nesting could not be confirmed among six spring and summer records. Age and/or sex determinations for 28 owls found the majority were females greater than one-year-old. Annual and seasonal chronologies derived from the records provide a more definitive status than previously available in the literature. The Long-eared Owl is a poorly understood species with little known about it regionally. This study provides some valuable insights to their life history on the Delmarva Peninsula furnishing baseline information for future studies.

ACKNOWLEDGMENTS

Personal observations, banding records, and/or databases contributed by Harry T. Armistead (Maryland Ornithological Society [MOS]), David F. Brinker (Central Regional Ecologist, Wildlife and Heritage Service, Maryland Department of Natural Resources, Annapolis, MD), Edward S. Brinkley (Editor, *North American Birds*, Cape Charles, VA), Katharine E. Duffy (Cape May Owl Banding Project [part of the Cape May Raptor Banding Project]), Robert F. Ringler (MOS), and Fletcher M. Smith (Research Biologist, The Center for Conservation Biology, College of William and Mary and Virginia Commonwealth University, Williamsburg, VA) added many records to the dataset. Danny Bystrak (Wildlife Biologist, Bird Banding Laboratory, Patuxent Wildlife Research Center, Laurel, MD) provided banding records from the USGS Bird Banding Laboratory database. David Bridge (MOS) and Jean L. Woods (Curator of Birds, Delaware Museum of Natural History, Wilmington, DE) located and confirmed data on museum specimens and provided copies of difficult to locate publications. Walter G. Ellison (Biology Lecturer, Washington College, Chestertown, MD), Jean L. Woods, and two anonymous reviewers made helpful comments on an earlier draft.

AUTHORS' APPEAL: Long-eared Owls are highly sensitive to disturbance. If you happen upon a perched owl, immediately exit the area, do not return that season, and do not relate the owl's specific location to anyone. You could publicly report the owl occurrence, but never give the specific location.

LITERATURE CITED

- Beardslee, C.S., and H.D. Mitchell. 1965. Birds of the Niagara Frontier Region: An annotated check-list. *Bulletin of the Buffalo Society of Natural Sciences* 22. 478 pp.
- Beaton, G., P.W. Sykes, Jr., and J.W. Parrish, Jr. 2003. *Annotated Checklist of Georgia Birds*. Occasional Publication No.14 (fifth edition). Georgia Ornithological Society. 156 pp.
- Bosakowski, T., R. Kane, and D.G. Smith. 1989. Decline of the Long-eared Owl in New Jersey. *The Wilson Bulletin* 101(3):481–485.
- Cheveau, M., P. Drapeau, L. Imbeau, and Y. Bergeron. 2004. Owl winter irruptions as an indicator of small mammal population cycles in the boreal forest of eastern North America. *Oikos* 107(1):190–198.
- Clark, R.J. 1975. A field study of the Short-eared Owl, *Asio flammeus* (Pontoppidan), in North America. *Wildlife Monographs* 47:1–67.
- Collins, C.T. 1962. Red-tailed Hawk attacks Long-eared Owl. *The Wilson Bulletin* 74(1):89.
- Delaware Division of Fish and Wildlife. 2015. 2015 Delaware Wildlife Action Plan. Division of Fish and Wildlife, Delaware Department of Natural Resources and Environmental Control. Available at: <http://www.dnrec.delaware.gov/fw/dwap/>. Accessed 27 January 2017.
- Duffy, K., and P. Kerlinger. 1992. Autumn owl migration at Cape May Point, New Jersey. *The Wilson Bulletin* 104(2):312–320.
- Ellison, W.G. 2010. Long-eared Owl, *Asio otus*. Pages 186–187 in: Ellison, W.G. (Editor), *2nd Atlas of the Breeding Birds of Maryland and the District of Columbia*. The Johns Hopkins University Press, Baltimore, MD. 494 pp.
- Environmental Systems Research Institute. 2012. ArcGIS 10.1. Available at: <http://www.esri.com>. Accessed 15 January 2016.
- Fletcher, A.J., R.B. Fletcher, M.W. Hewitt, A.O’C. Knotts, and A.M. Thompson. 1956. *List of Caroline County Birds*. Maryland Avifauna Number 1, Maryland Ornithological Society, Inc., Baltimore, MD. 22 pp.
- Fletcher, A.J., R.B. Fletcher, M.W. Hewitt, A.O’C. Knotts, A.M. Thompson, and V.E. Unger. 1979. *List of Caroline County Birds* (Revised 31 January

- 1979). Maryland Avifauna Number 1, Maryland Ornithological Society, Inc., Baltimore, MD. 29 pp.
- Gross, D.A. 2012. Long-eared Owl, *Asio otus*. Pages 210–211 in: Wilson, A.M., D.W. Brauning, and R.S. Mulvihill (Editors), *Second Atlas of Breeding Birds in Pennsylvania*. The Pennsylvania State University Press, University Park, PA. 586 pp.
- Hampe, I.E., and H. Kolb. 1947. *A Preliminary List of Birds of Maryland and the District of Columbia*. The Natural History Society of Maryland, Baltimore, MD. 80 pp.
- Harvey, P.V., and N. Riddiford. 1990. An uneven sex ratio of migrant Long-eared Owls. *Ring and Migration* 11(3):132–135.
- Hess, G.K., R.L. West, M.V. Barnhill III, and L.M. Fleming. 2000. *Birds of Delaware*. University of Pittsburgh Press, Pittsburgh, PA. 635 pp.
- Kain, T. (Editor). 1987. *Virginia's Birdlife: An annotated checklist*. Virginia Society of Ornithology. Virginia Avifauna Number 3. 127 pp.
- Kirkwood, F.C. 1895. A list of the birds of Maryland. *Transactions of the Maryland Academy of Science* 2:241–382.
- Kolb, H. 1947a. Breeding of the Long-eared Owl near Baltimore. *Maryland: A Journal of Natural History* 17(2):23–25.
- Kolb, H. 1947b. Breeding of the Long-eared Owl near Baltimore. *Maryland Conservationist* 24(2):18, 22.
- Korpimäki, E., and C.J. Krebs. 1996. Predation and population cycles of small mammals: A reassessment of the predation hypothesis. *BioScience* 46(10):754–764.
- Korpimäki, E., and K. Norrdahl. 1991. Numerical and functional responses of Kestrels, Short-eared Owls, and Long-eared Owls to vole densities. *Ecology* 72(3):814–826.
- Larner, Y.R. (Chairman of the Checklist Committee). 1979. *Virginia's Birdlife: An annotated checklist*. Virginia Society of Ornithology. Virginia Avifauna Number 2. 118 pp.
- Marks, J.S., D.L. Evans, and D.W. Holt. 1994. Long-eared Owl, *Asio otus*. The Birds of North America (P. G. Rodewald, Editor). Cornell Lab of

Ornithology, Ithaca, NY. Available at: <https://birdsna.org/Species-Account/bna/species/loewowl>. Accessed 1 March 2016.

Maryland Department of Natural Resources. 2015. 2015 Maryland State Wildlife Action Plan. Wildlife and Heritage Service, Maryland Department of Natural Resources. Available at: http://dnr2.maryland.gov/wildlife/Pages/plants_wildlife/SWAP_Submission.aspx. Accessed 27 January 2017.

Mississippi Ornithological Society. 2015. Checklist of Birds of Mississippi. Mississippi Ornithological Society Bird Records Committee. 2 pp. Available at: http://www.missbird.org/Files/Mississippi%20State%20Checklist/MOS_Checklist_Aug_2015.pdf. Accessed 27 January 2017.

Murray, J.J. (Editor). 1952. *A Check-list of the Birds of Virginia*. Virginia Society of Ornithology. 113 pp.

Overskaug, K., and E. Kristiansen. 1994. Sex ratio of accidentally killed Long-eared Owls *Asio otus* in Norway. *Ringing and Migration* 15(2):104–106.

Pennock, C.J. 1904. *Some of Our Useful Birds*. Bulletin No. 5. State Board of Agriculture, Dover, DE. 16 pp.

Pitelka, F.A., P.Q. Tomich, and G.W. Treichel. 1955a. Breeding behavior of jaegers and owls near Barrow, Alaska. *The Condor* 57(1):3–18.

Pitelka, F.A., P.Q. Tomich, and G.W. Treichel. 1955b. Ecological relations of jaegers and owls as lemming predators near Barrow, Alaska. *Ecological Monographs* 25(1):85–117.

Potter, J.K. 1937. The Season, CXVII April 15 to June 15, 1937: Philadelphia Region. *Bird Lore* 39(4):312–326.

Pyle, P. 1997. *Identification Guide to North American Birds, Part I: Columbidae to Ploceidae*. Slate Creek Press, Bolinas, CA. 732 pp.

R Development Core Team. 2016. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. (software)

Rhoads, S.N., and C.J. Pennock. 1905. Birds of Delaware: A preliminary list. *The Auk* 22(2):194–205.

- Rives, W.C. 1890. *A Catalogue of the Birds of the Virginias*. Proceedings of the Newport Natural History Society 1889–90, Document VII. Newport, RI. 100 pp.
- Rottenborn, S.C., and E.S. Brinkley. 2007. *Virginia's Birdlife: An annotated checklist*. Virginia Avifauna, No. 7 (Fourth Edition), Virginia Society of Ornithology. 330 pp.
- RStudio. 2015. RStudio: Integrated Development Environment for R (Version 0.99.491). RStudio, Boston, MA. (software)
- Russell, R.W., P. Dunne, C. Sutton, and P. Kerlinger. 1991. A visual study of migrating owls at Cape May Point, New Jersey. *The Condor* 93(1):55–61.
- Santner, S. 1992. Long-eared Owl, *Asio otus*. Pages 162–163 in: Brauning, D.W. (Editor). *Atlas of Breeding Birds in Pennsylvania*. University of Pittsburgh Press, Pittsburgh, PA. 484 pp.
- Slack, R.S., C.B. Slack, R.N. Roberts, and D.E. Emord. 1987. Spring migration of Long-eared Owls and Northern Saw-whet Owls at Nine Mile Point, New York. *The Wilson Bulletin* 99(3):480–485.
- Stewart, R.E., and C.S. Robbins. 1958. *Birds of Maryland and the District of Columbia*. North American Fauna, No. 62, United States Government Printing Office, Washington D.C. 401 pp.
- Stone, W.T. 1965. *Bird studies at Old Cape May: An ornithology of Coastal New Jersey*, Volume II (reprint). Dover Publications, Inc., New York, NY. 457 pp.
- Village, A. 1981. The diet and breeding of Long-eared Owls in relation to vole numbers. *Bird Study* 28(3):215–224.
- Village, A. 1987. Numbers, territory-size and turnover of Short-eared Owls *Asio flammeus* in relation to vole abundance. *Ornis Scandinavica* 18(3):198–204.
- Virginia Department of Game and Inland Fisheries. 2015. Virginia's 2015 Wildlife Action Plan. Available at: <http://bewildvirginia.org/wildlife-action-plan/>. Accessed 27 January 2017.
- Wyllie, I., L. Dale, and I. Newton. 1996. Unequal sex-ratio, mortality causes and pollutant residues in Long-eared Owls in Britain. *British Birds* 89(10):429–436.

Novel Winter Bathing Behavior by a Merlin (*Falco columbarius*)

David R. Smith

2203 Flag Marsh Road, Mount Airy, Maryland 21771
Lacsmith12@comcast.net

Few reports of bathing in the family Falconidae have been published. Bent (1938) mentions it only for the Prairie Falcon (*Falco mexicanus*), as does Holthuijzen et al. (1987). Palmer (1988) describes bathing behavior for the Peregrine Falcon (*F. peregrinus*). Ristow et al. (1980) describes bathing by Eleonora's Falcon (*F. eleonora*), as does Eisermann (2005) for the Orange-breasted Falcon (*F. deiroleucus*). For the Merlin (*F. columbarius*), Warkentin et al. (2005) reported one bathing in rainwater, and Haak and Buchanan (2012) reported several instances of winter bathing behavior. Most of these accounts are of birds wading in shallow water.

Schmidl (1988) lists various observed bathing techniques by falcons, including stand-in bathing in shallow water, flight bathing on the wing through a series of dips and rises, rain bathing either perched or on the wing, and snow bathing. Palmer (1988) also describes flight-bathing behavior, whereby a flying falcon briefly contacts the surface of a water body at a low angle before immediately ascending. Eisermann (2005) reported a different type of flight-bathing behavior whereby Orange-breasted Falcons flew directly into rain-soaked leaves. Swann et al. (1993) report several instances of immature Peregrine Falcons engaging in swimming behavior, and speculate that the behavior was associated with bathing. In this account, I report on a presumed bathing behavior of a Merlin using a swimming technique.

OBSERVATION

The observation occurred from the Chesapeake & Ohio Canal National Historical Park (C&O Canal) towpath. The location of the observation was upriver from Milepost 91 near the confluence of Opequon Creek with the Potomac River. Temperatures at the start of the morning had been around (-9° C [15° F]) but by the time of the observation were just above freezing (0° C [32° F]). The observation was made using a 10x Zeiss binocular.

At approximately 1140 hours on 30 January 2016, I observed a Merlin flying several meters above the iced-over Potomac River. The falcon flew near the West Virginia shoreline within 200 m (656 ft) of where I stood, swooped low

over the river where the open Opequon Creek met the frozen Potomac River, then landed on the West Virginia shoreline on a broken tree trunk. Within about one minute, the Merlin flew from its perch and landed on the edge of the ice where the open water of Opequon Creek met the Potomac River. The Merlin stood in this location for another minute and then jumped into the open water breast first. It immediately flew from the water and landed at the edge of the ice. There it remained relatively motionless for several minutes, scanning the area by turning its head from side to side. The Merlin again suddenly plunged into the open water breast first, but rather than exiting the water immediately, it began to use its wings to row into open water away from the edge of the ice. It rowed in a shallow circle approximately 2 m (~7 ft) away from the ice shelf, returned to the edge of the ice, flapped out of the water, and stood on the ice. It then shook its feathers and soon flew back to the previous perch on the broken tree trunk. The Merlin then proceeded to preen its feathers. The entire sequence of activities lasted approximately 10 minutes, after which I departed with the Merlin still perched.

DISCUSSION

While few reported observations exist of bathing Merlins and none that I could find that involved swimming, it is well known that falcons regularly bathe. For Peregrine Falcons, Palmer (1988) suggested that this has been known for centuries. Swann et al. (1993) also stated that, "Peregrines have a strong 'motivation' to bathe daily." At a falconry event in Utah during late autumn, falcon owners, including those with Merlins, would break ice in frozen bird baths to allow their birds to bathe (Tom Humphrey, pers. comm., 4 February 2016). The observation of a Merlin bathing in winter by swimming may have been a result of a scarcity of open water resulting from extremely cold temperatures in the area at the time and an opportunity to exploit access to open water in an area with good visibility of the surroundings, which enabled scanning for potential predators. Swann et al. (1993) postulated a similar explanation for immature Peregrine Falcons swimming in Maine where access to shallow water for wading was limited.

ACKNOWLEDGMENTS

I thank Tom Humphrey (falconer and immediate Past President of the Frederick Bird Club) for relating the story to me about falconers breaking ice to allow their falcons to bathe. I also thank Joseph B. Buchanan (Natural Resource Scientist, Washington Department of Fish and Wildlife, Olympia, WA) for his thoughtful review of an earlier draft and for providing additional falcon bathing references. Comments of two other anonymous reviewers also improved an earlier draft.

LITERATURE CITED

- Bent, A.C. 1938. *Life Histories of North American Birds of Prey, (Part 2: Orders Falconiformes and Strigiformes)*. Smithsonian Institution, United States National Museum, Bulletin 170. Washington, DC. 482 pp.
- Eisermann, K. 2005. An observation of foliage-bathing by an Orange-breasted Falcon (*Falco deiroleucus*) in Tikal, Guatemala. *The Wilson Bulletin* 117(4):415–418.
- Haak, B.A., and J.B. Buchanan. 2012. Bathing and drinking behavior of wintering Merlins. *Journal of Raptor Research* 46(2):224–226.
- Holthuijzen, A.M.A., P.A. Duley, J.C. Hagar, S.A. Smith, and K.N. Wood. 1987. Bathing behavior of nesting Prairie Falcons (*Falco mexicanus*) in southwestern Idaho. *The Wilson Bulletin* 99(1):135–136.
- Palmer, R.S. (Editor). 1988. *Handbook of North American Birds, Volume 5: Diurnal Raptors (Part 2)*. Yale University Press, New Haven, CT. 465 pp.
- Ristow, D., C. Wink, and M. Wink. 1980. The bathing behaviour of Eleonora's Falcon. *Bird Study* 27(1):54–56.
- Schmidl, D. 1988. Dusting in falcons. *Journal of Raptor Research* 22(2):59–61.
- Swann, D., T. Mahoney, and R.D. Applegate. 1993. Swimming behavior by immature Peregrine Falcons. *Maine Naturalist* 1(1):36–37.
- Warkentin, I.G., N.S. Sodhi, R.H.M. Espie, A.F. Poole, L.W. Oliphant, and P.C. James. 2005. Merlin (*Falco columbarius*), *The Birds of North America* (P.G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca NY. Available at: <https://birdsna.org/Species-Account/bna/species/merlin>. Accessed 16 February 2017.

First Maryland Record of Pacific-slope/Cordilleran (“Western”) Flycatcher, *Empidonax difficilis/occidentalis*

Kathleen B. Calvert

2404 Doubs Court, Adamstown, Maryland 21710
kathycalvert1@gmail.com

On 16 November 2015, I photographed an *Empidonax* flycatcher in southwestern Caroline County, Maryland. The bird caught my eye as I was birding along Poplar Neck Road at the edge of a scrubby field on the northeast side of the road. I had been playing a tape of Eastern Screech-Owl, *Megascops asio*, in order to attract the attention of passerines or other birds that might be in the area. At 11:55 a.m., out of the corner of my eye, I noted a bird actively flycatching in the small deciduous trees at the edge of the field about 50 ft (15 m) from where I was positioned. Although it is possible that the recording attracted the bird into the general area, it did not seem particularly interested in the sound from the tape. The bird was very active, perching on a branch then suddenly sallying out for an insect and flying back to the same perch or one nearby. Each time, when it first perched, the bird flicked its tail several times. It continued flycatching, sometimes landing in sight, but more often perching on branches that were out of my line of sight a few yards deeper inside the edge of trees and bushes. The bird seemed to do most of its foraging at about 3–5 ft (1–1.5 m) above the ground.

The bird was clearly an *Empidonax* (empid), based on its small size, olive upperparts, pale underparts, wing bars, and eye ring. Any empid is an unexpected species in late November in Maryland and is likely to be a rare species, or a very late-date breeding or migratory species. I noted that this bird had very bright yellow underparts, with a hint of olive on the upper breast, and a pale throat (Figure 1). The bird had a very prominent pale eye ring that was narrower at the top and bottom and extended out in a teardrop fashion at the rear. My photos did not capture the hint of a peak to the back of the head, which I saw through binoculars. Its wings were a sooty dark color with two light buff-colored wing-bars (Figure 2). I was never able to get a good sense of the relative length of the tail, but the primaries did not seem to extend to the base of the tail. The bird did not vocalize.

I made phone calls and sent texts with photos immediately, but unfortunately the bird was not seen again that day or in following days despite multiple efforts by other birders to re-find it. However, my description, photos, and the timing of



Figure 1. Pacific-slope/Cordilleran Flycatcher (“Western”) Flycatcher, *Empidonax difficilis/occidentalis*. Poplar Neck Road near Choptank, Caroline County, Maryland, 16 November 2015.



Figure 2. Pacific-slope/Cordilleran Flycatcher (“Western”) Flycatcher. Poplar Neck Road near Choptank, Caroline County, Maryland, 16 November 2015.

the bird's occurrence were sufficient to rule out Yellow-bellied Flycatcher (*Empidonax flaviventris*), the only other species with which this bird might have been confused. The Yellow-bellied Flycatcher is typically gone from Maryland by mid-October. The eye ring of the Yellow-bellied Flycatcher is usually circular and more rounded behind the eye, not showing the distinct teardrop profile of the bird I saw. The coloration of the underparts of the bird on Poplar Neck Road was also much brighter yellow than would be expected on a Yellow-bellied Flycatcher. Regarding the bill, which did not appear especially wide, I noted that the upper mandible was dark and the lower was yellowish-orange. The bill seemed a bit longer than what I would have expected for a Yellow-bellied Flycatcher, although Yellow-bellied does show a lot of variation in bill shape and size. Analysis of the secondaries (Figure 2) showed a buff-colored fringing pattern consistent with Pacific-slope/Cordilleran ("Western") Flycatcher (*Empidonax difficilis/occidentalis*) as compared to Yellow-bellied Flycatcher (Baumann et al. 2014). I submitted my photos and a written report to the Maryland/DC Records Committee, and the record was accepted in 2016 as a Pacific-slope/Cordilleran Flycatcher.

THE "WESTERN" FLYCATCHER SPECIES COMPLEX

In 1989, The American Ornithologists' Union split Western Flycatcher (*Empidonax difficilis*) into two species: Pacific-slope Flycatcher (*E. difficilis*) and Cordilleran Flycatcher (*E. occidentalis*) based on the work of Johnson and Marten (1988). The normal range for Pacific-slope Flycatcher is the Pacific coastal mountain ranges of Canada, the United States, and Mexico (Lowther et al. 2016b); for the Cordilleran Flycatcher, the range is the Rocky Mountains of the United States and Mexico (Lowther et al. 2016a). The two species are virtually indistinguishable in the wild (Lowther et al. 2016a, 2016b). In the western United States, the best characters for separation are voice, range, and habitat (Lowther et al. 2016a, 2016b). Obviously range and habitat are of no value when trying to identify these rare Northeast and Atlantic Coast vagrants. Voice is not always usable in the East, but DNA of scat (i.e., environmental DNA or eDNA) has occasionally been used to separate the species (eBird 2017).

SUMMARY OF NORTHEAST AND ATLANTIC COAST RECORDS

Members of the "Western" Flycatcher species complex are rare in the Northeast and along the Atlantic Coast. Table 1 shows the twelve state-bird-records-committee-accepted records for the species complex and whether the record was accepted for Pacific-slope or Pacific-slope/Cordilleran Flycatcher. There are no eastern records for Cordilleran Flycatcher. Only records-committee-accepted records are included in this list. The records cluster geographically between Massachusetts and North Carolina, with two outliers in Florida. Records have occurred between 14 September and 15 January, with five occurring in

November and three in December. The earliest accepted record occurred in 1981. When the decades are compared, one record occurred in the 1980s, three in the 1990s, three in the 2000s, and five in the 2010s.

Table 1. State-bird-records-committee-accepted records of Pacific-slope/Cordilleran Flycatcher for the Northeast and Atlantic Coast states.

| State | Date | Location | Taxon | Citation |
|----------------|------------------------------|--|-------------------------------|---|
| Maine | (no records) | | | MBRC 2017 |
| Vermont | (no records) | | | VBRC 2016 |
| New Hampshire | (no records) | | | NHRBC 2010 |
| Massachusetts | 10 Nov 2006 | Manomet, Plymouth, Plymouth Co. | Pacific-slope/ Cordilleran | MARC 2013 |
| Rhode Island | (no records) | | | RIARC 2016 |
| Connecticut | 1–7 Dec 2007 | Osbornedale State Park, Derby, New Haven Co. | Pacific-slope/ Cordilleran | ARCC 2016 |
| Connecticut | 19–20 Dec 2015 | Branford, New Haven Co. | Pacific-slope/ Cordilleran | ARCC 2016 |
| New York | 14–16 Sep 1995 | Fire Island Lighthouse, Suffolk Co. | Pacific-slope/ Cordilleran | NYSARC 2015 |
| Pennsylvania | 16 Dec 1990 | East Drumore Township, Lancaster Co. | Pacific-slope | PORC 2017 |
| Pennsylvania | 23 Nov 2012 | State Game Lands 230, Cumberland Co. | Pacific-slope | PORC 2017 |
| New Jersey | 16 Nov 1981 | Brigantine NWR, Atlantic Co. | Pacific-slope/ Cordilleran | NJBRC 2016 |
| Maryland/DC | 16 Nov 2015 (this record) | Poplar Neck Rd., Choptank, Caroline Co. | Pacific-slope/ Cordilleran | MD/DCRC 2016 |
| Delaware | (no records) | | | DBRC 2015 |
| West Virginia | (no records) | | | WVBRC 2016 |
| Virginia | 12 Nov 1993 | Eastern Shore of Virginia NWR, Northampton Co. | Pacific-slope/ Cordilleran | VARCOM 2017 |
| North Carolina | 15 Jan 2000 | Jordan Lake, Chatham Co. | Pacific-slope/ Cordilleran | CBCBRC 2015 |
| South Carolina | (no records) | | | CBCBRC 2015 |
| Georgia | (no records) | | | GCRC 2015 |
| Florida | 8 Jan 2015 | Manakin Manor South, Leon Co. | Pacific-slope/ Cordilleran | FOSRC 2013, Greenlaw 2015 |
| Florida | 1–2 October 2015 | Fort Pickens, Escambia Co. | Pacific-slope/ Cordilleran | Andrew W. Kratter in litt. 28 Mar 2017 |

Table 2 summarizes reports submitted to eBird (not necessarily officially-accepted records) of eastern sightings for Pacific-slope or Pacific-slope/Cordilleran Flycatcher. There are no Cordilleran Flycatcher reports. Some of these reports are merely date extensions of the Table 1 officially-accepted

records. Table 2 adds additional reports not found in Table 1 for New York, Pennsylvania, and Florida. Again, the reports cluster geographically between Massachusetts and North Carolina, with one outlier in Florida. Reports have occurred between 31 October and 22 January, with the majority of reports occurring from November to January.

Table 2. Submitted eBird Reports (not necessarily official state-bird-records-committee-accepted records) of Pacific-slope/Cordilleran Flycatcher for the Northeast and Atlantic Coast states (eBird 2017). “P-s/C” = Pacific-slope/Cordilleran

| State | Date | Location | Submitted Taxon | State-accepted Record |
|----------------|---------------------------|--|----------------------------|--|
| Maine | (no reports) | | | |
| Vermont | (no reports) | | | |
| New Hampshire | (no reports) | | | |
| Massachusetts | 10 Nov 2006 | Manomet Center for Conservation Sciences, Plymouth, Plymouth Co. | Pacific-slope/ Cordilleran | yes |
| Rhode Island | (no reports) | | | |
| Connecticut | 19–20 Dec 2015 | Branford Hills School, Burban Dr., Branford, New Haven Co. | Pacific-slope | yes, as P-s/C |
| New York | 18–23 Nov 2015 | Central Park, New York, New York Co. | Pacific-slope | for future review (Willie D’Anna, in litt., 26 Mar 2017) |
| New York | 8–10 Dec 2016 | Inwood Hill Park, New York, New York Co. | Pacific-slope/ Cordilleran | for future review (Willie D’Anna, in litt., 26 Mar 2017) |
| Pennsylvania | 18 Dec 1990 | East Drumore Township, Lancaster Co. | Pacific-slope | date extension |
| Pennsylvania | 18 Nov 2012 | State Game Lands 230, Cumberland Co. | Pacific-slope/ Cordilleran | date extension as P-s |
| Pennsylvania | 24–25 Nov 2012 | State Game Lands 230, Cumberland Co. | Pacific-slope | date extension |
| Pennsylvania | 20 Dec 2015–5 Jan 2016 | Berks County Heritage Center, Berks Co. | Pacific-slope | submitted to PORC for review |
| New Jersey | (no reports) | | | |
| Maryland/DC | 16 Nov 2015 (this record) | Poplar Neck Rd., Choptank, Caroline Co. | Pacific-slope/ Cordilleran | yes |
| Delaware | (no reports) | | | |
| West Virginia | (no reports) | | | |
| Virginia | 12 Nov 1993 | Eastern Shore of Virginia NWR, Northampton Co. | Pacific-slope | yes, as P-s/C |
| Virginia | 29 Nov 1993 | Eastern Shore of Virginia NWR, Northampton Co. | Pacific-slope | date extension? |
| North Carolina | 15–22 Jan 2000 | Jordan Lake, Chatham County | Pacific-slope/ Cordilleran | date extension |
| South Carolina | (no reports) | | | |
| Georgia | (no reports) | | | |
| Florida | 31 Oct 2004 | Fort De Soto Park, Pinellas Co. | Pacific-slope/ Cordilleran | not submitted to FOSRC (Andrew W. Kratter in litt., 28 Mar 2017) |

ACKNOWLEDGMENTS

I thank Eugene J. Scarpulla for assistance locating “Western” Flycatcher records, as well as Willie D’Anna (New York State Avian Records Committee) and Andrew W. Kratter (Florida Ornithological Society Records Committee) for updated information from their respective state records committees. I also thank two anonymous reviewers.

LITERATURE CITED

- AOU (American Ornithologists’ Union). 1989. Thirty-Seventh Supplement to the American Ornithologists’ Union Check-List of North American Birds. *The Auk* 106(3):532–538.
- ARCC (Avian Records Committee of Connecticut). 2016. Species Index by Common Name for the Published Reports of ARCC. Available at: http://www.ctbirding.org/ARCC/index_of_reports_by_species.htm. Accessed 30 January 2017.
- Baumann, M.J., S.C. Galen, N.D. Pederson, and C.C. Witt. 2014. Simple technique for distinguishing Yellow-bellied Flycatchers from Cordilleran and Pacific-slope Flycatchers. *Journal of Field Ornithology* 85(4):391–396.
- CBCBRC (Carolina Bird Club Bird Records Committees). 2015. *Chat* databases. Available at: <https://www.carolinabirdclub.org/brc/>. Accessed 4 February 2017.
- DBRC (Delaware Bird Records Committee). 2015. Summary of Completed Reviews (1 September 2015). Available at: <http://www.dosbirds.org/committees/records/>. Accessed 4 February 2017.
- eBird. 2017. Pacific-slope Flycatcher, Cordilleran Flycatcher, and Pacific-slope/Cordilleran Flycatcher (Western Flycatcher). eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available at: <http://www.ebird.org>. Accessed 5 February 2017.
- FOSRC (Florida Ornithological Society Records Committee). 2013. Official Florida State Bird List (updated through 31 August 2013). Available at: <http://www.fosbirds.org/content/records-committee>. Accessed 4 February 2017.

GCRC (Georgia Ornithological Society Checklist and Records Committee).

2015. 2015 Checklist of Georgia Birds. Available at: <http://gos.org/checklist-and-records-committee>. Accessed 4 February 2017.

Greenlaw, J.S. FOSRC Records Committee Board Report. 8 August 2015

FOSRC (Florida Ornithological Society Records Committee) Meeting.

Available at: http://www.fosbirds.org/sites/default/files/fosrc_report_fall_2015_leesburg.pdf. Accessed 25 January 2016.

Johnson, N.K., and J.A. Marten. 1988. Evolutionary genetics of flycatchers. II.

Differentiation in the *Empidonax difficilis* complex. *The Auk* 105(1):177–191.

Lowther, P.E., P. Pyle, and M.A. Patten. 2016a. Cordilleran Flycatcher

(*Empidonax occidentalis*), The Birds of North America (P.G. Rodewald,

Editor). Cornell Lab of Ornithology, Ithaca, NY. Available at:

<https://birdsna.org/Species-Account/bna/species/corfly>. DOI:

10.2173/bna.556b. Accessed 5 February 2017.

Lowther, P.E., P. Pyle, and M.A. Patten. 2016b. Pacific-slope Flycatcher

(*Empidonax difficilis*), The Birds of North America (P.G. Rodewald, Editor).

Cornell Lab of Ornithology, Ithaca, NY. Available at:

<https://birdsna.org/Species-Account/bna/species/pasfly>. DOI:

10.2173/bna.556a. Accessed 5 February 2017.

MARC (Massachusetts Avian Records Committee). 2013. Official

Massachusetts State List (March 2013). Available at:

<http://www.maavianrecords.com/>. Accessed 4 February 2017.

MBRC (Maine Bird Records Committee). 2017. Official List of Maine Birds

(updated 1 January 2017). Available at:

<https://sites.google.com/site/mainebirdrecordscommittee/>. Accessed 30

January 2017.

MD/DCRC (Maryland/District of Columbia Records Committee). 2016.

MD/DCRC Database Abridged Version, Maryland, 7 November 2016.

Available at: <http://www.mdbirds.org/mddcrc/pdf/mddatabase.pdf>. Accessed 30 January 2017.

NHRBC (New Hampshire Rare Bird Committee). 2010. The New Hampshire

State Bird List (20 April 2010). Available at: <http://nhbirdrecords.org/>.

Accessed 30 January 2017.

- NJBRC (New Jersey Bird Records Committee). 2016. New Jersey Bird Records Committee List of Accepted Records of Rare Birds in New Jersey through December 2015 (seventh corrected and revised edition, August 2016). Available at: <http://www.njbrc.net/>. Accessed 4 February 2017.
- NYSARC (New York State Avian Records Committee). 2015. NYSARC Reports and Decisions through 2012 (updated 18 November 2015). Available at: <http://nybirds.org/NYSARC/index.htm>. Accessed 4 February 2017.
- PORC (Pennsylvania Ornithological Records Committee). 2017. Database, and Search. Available at: <http://www.pabirds.org/records/>. Accessed 26 March 2017.
- RIARC (Rhode Island Avian Records Committee). 2016. Checklist of Rhode Island Birds (July 2016). Available at: <http://www.ri-avianrecords.org/>. Accessed 30 January 2017.
- VARCOM (Virginia Avian Records Committee). 2017. Database for Ornithological Verification and Submission (DOVES). Available at: <http://www.virginiabirds.net/varcom.html>. Accessed 4 February 2017.
- VBRC (Vermont Bird Records Committee). 2016. Vermont State Checklist (updated 15 November 2016). Available at: <http://vtcostudies.org/wildlife/wildlife-watching/vbrc/>. Accessed 30 January 2017.
- WVBRC (West Virginia Bird Records Committee). 2016. West Virginia State Bird List (5 March 2016). Available at: <http://www.brooksbirdclub.org/west-virginia-bird-records-committee.html>. Accessed 4 February 2017.

Sunbathing by Great Crested Flycatchers, *Myiarchus crinitus*

Barbara K. Johnson

1905 Kingswood Court, Annapolis, Maryland 21401
barbarajohnson222@gmail.com

Beginning summer 2008, I observed Great Crested Flycatchers, *Myiarchus crinitus*, each breeding season at the above address, on a lot dominated by mature American beech (*Fagus grandifolia* Ehrh.), tulip poplar (*Liriodendron tulipifera* L.), and several oak species (*Quercus* L. spp.). In 2008 and 2009, a pair nested nearly 50 ft (15 m) from the house, in a visible tree cavity. On numerous occasions between July 2009 and August 2016, I observed a Great Crested Flycatcher sunbathing on the vinyl cover of an outdoor spa on the house's deck. On two occasions, I recorded two Great Crested Flycatchers engaged in the behavior simultaneously, and on several occasions, one flycatcher was seen sunbathing while another was visible or calling nearby.

In 2015 and 2016, I recorded several sunbathings, and for some, but not all, I noted the date, time of day, air temperature, relative humidity, temperature of the vinyl surface (using an infrared or laser thermometer), duration of sunbathing, and the birds' activity. Sixteen sunbathing incidents were recorded on eight dates; in 2015: 6, 10, 19, 20, and 28 July, and 3 August; and in 2016: 11 and 16 July (Table 1). On the latter date, the accompanying photos were taken during multiple visits (Figures 1–4).

All observations began between 11:50 a.m. and 5:35 p.m. Duration of contact with the surface ranged from approximately 10 seconds to approximately 150 seconds. Air temperature ranged from 80° F (26.7° C) to 90° F (32.2° C). Relative humidity ranged from 39% to 71%.

The temperature of the vinyl surface (measured within several seconds of the birds' departure) ranged from 135.2° F (57.3° C) to 182.4° F (83.6° C). The vinyl surface temperature, unpleasantly hot to the touch, was higher than that of the surrounding wood deck by as much as 51° F (10.6° C).

Typical sunbathing activity included a bird landing first in a nearby tree, then flying down to the vinyl and immediately splaying wings and tail against the surface, then either scooting (propelling its body across the vinyl, its ventral surface constantly pressed against the vinyl, covering distances ranging from approximately 2–3 in [5.0–7.6 cm] up to approximately 6 ft [1.8 m]), or

remaining still, or alternating between scooting and stillness. Often, the head was cocked upward and the mouth gaped (once for approximately 120 seconds), most often while the bird remained still. When the bird scooted, it would maneuver so as to expose varying areas of its ventral surface to the vinyl. The longest observed period of scooting was approximately 17 seconds, and the longest stationary period was approximately 140 seconds.

Table 1. Incidental data collection of Great Crested Flycatcher sunbathing behavior, 2015 and 2016. (□ = no data collected)

| Date | Time | Air | Relative | Surface | Duration (seconds) | Scooting (seconds) | Stationary (seconds) |
|-------------|----------|--------------------------|-----------------|--------------------------|-----------------------|-----------------------|-------------------------|
| | | Temperature (° F/° C) | Humidity (%) | Temperature (° F/° C) | | | |
| 06 Jul 2015 | 1600 | □ | □ | □ | □ | □ | □ |
| 10 Jul 2015 | 1150 | 80/26.7 | 48 | □ | □ | □ | □ |
| 10 Jul 2015 | visit #2 | □ | □ | □ | □ | □ | □ |
| 19 Jul 2015 | 1201 | 90/32.2 | 71 | 173.0/78.3 | 150 | 10 | 140 |
| 20 Jul 2015 | 1343 | 90/32.2 | 57 | 182.4/83.6 | 21 | 17 | 4 |
| 28 Jul 2015 | 1543 | 89/31.7 | 50 | 158.4/70.2 | 140 | 10 | 130 |
| 28 Jul 2015 | 1548 | 89/31.7 | 50 | 159.2/70.7 | 106 | 4 | 102 |
| 03 Aug 2015 | 1250 | 88/31.1 | 39 | 181.0/82.8 | 10 | 10 | 0 |
| 11 Jul 2016 | 1630 | □ | □ | 135.2/57.3 | □ | □ | □ |
| 11 Jul 2016 | visit #2 | □ | □ | □ | □ | □ | □ |
| 11 Jul 2016 | visit #3 | □ | □ | □ | □ | □ | □ |
| 11 Jul 2016 | 1735 | □ | □ | □ | □ | □ | □ |
| 16 Jul 2016 | 1555 | 86/30.0 | □ | 163.4/73.0 | 104 | 0 | 104 |
| 16 Jul 2016 | 1600 | □ | □ | □ | 60 | □ | □ |
| 16 Jul 2016 | 1610 | □ | □ | □ | 60 | □ | □ |
| 16 Jul 2016 | 1625 | □ | □ | □ | 92 | □ | □ |

On 11 July 2016, four sunbathings were recorded between 4:30 p.m. and 5:35 p.m. On 16 July 2016, four sunbathings were recorded between 3:55 p.m. and 4:25 p.m.

While two other species—Carolina Wren (*Thryothorus ludovicianus*) and Blue Jay (*Cyanocitta cristata*)—were observed sunbathing on the wooden deck and rail, only the Great Crested Flycatchers were seen sunbathing on the vinyl. The Great Cresteds were never observed sunbathing on any surface other than the vinyl.

DISCUSSION

Published literature reports sunbathing by various species of birds. Hauser (1957) proposed two categories of sunbathing: “voluntary,” where a bird



Figure 1. Great Crested Flycatcher sunbathing location on the vinyl cover of an outdoor spa. Annapolis, Maryland, 16 July 2016. (Note that a Blue Jay is also sunbathing on the railing)



Figure 2. Great Crested Flycatcher sunbathing. Facing right. Annapolis, Maryland, 16 July 2016.




Figure 3. Great Crested Flycatcher sunbathing. Facing front. Annapolis, Maryland, 16 July 2016.




Figure 4. Great Crested Flycatcher sunbathing. Facing away. Annapolis, Maryland, 16 July 2016.

intentionally seeks a sunlit area, and “compulsory,” where the bird does not seek the situation (e.g., on a sunlit shiny metal platform feeder). Goodwin (1967) also mentioned shielding of nestlings by raptors and herons as a form of compulsory sunbathing, but stated that most sunbathing is voluntary. Kennedy (1969) summarized six possible functions for voluntary sunbathing: 1) sunrise thermoregulation as observed in Turkey Vultures (*Cathartes aura*), Black Vultures (*Coragyps atratus*), and Double-crested Cormorants (*Phalacrocorax auritus*) (Curry-Lindahl [1970] also documented this behavior in Reed Cormorants [*Microcarbo africanus*].); 2) ectoparasite control; 3) drying, as observed in cormorants and Anhingas (*Anhinga anhinga*) after leaving the water; 4) vitamin D production from a precursor in the uropygial gland secretion; 5) a possible role in molting; and 6) increased production of the uropygial gland secretion. Horsfall (1984) further elaborated on these possible functions. Lanyon (1958), studying a hand-raised Western Meadowlark (*Sturnella neglecta*) and a similarly-raised European Starling (*Sturnus vulgaris*), observed that sunbathing was initiated by heat and not light.

Although Miller and Lanyon’s review (2014) found no published data on sunbathing by Great Crested Flycatchers, Hauser (1957) had previously observed three Great Crested Flycatchers on 25 June 1955, 1:30–3:00 p.m., sunbathing on a primarily elm (*Ulmus* L. sp.) leaf compost heap. The air temperature was 90° F (32° C) and that of the compost heap’s surface was 140° F (60° C). My Internet search also found several photos of sunbathing by Great Crested Flycatchers (Brown 2011, Tallman 2011, Roberts 2015, Zee 2015, Bacquie 2016).

Five other *Myiarchus* species (Ash-throated Flycatcher, *M. cinerascens*; Brown-crested Flycatcher, *M. tyrannulus*; Dusky-capped Flycatcher, *M. tuberculifer*; La Sagra’s Flycatcher, *M. sagrae*; and Nutting’s Flycatcher, *M. nuttingi*) can occur in the United States (Dunn and Alderfer 2006). Cardiff and Dittmann (2000) reported that there was no published literature on sunbathing for Ash-throated Flycatcher. They reported the same for Brown-crested Flycatcher (Cardiff and Dittmann 2002). I could find no reports for those two species or for either La Sagra’s Flycatcher or Nutting’s Flycatcher. Tweit and Tweit (2002) reported that Skutch (1960) had observed sunbathing by a Dusky-capped Flycatcher. Skutch’s account stated:

“Once while I sat in a blind in the forest on Barro Colorado Island [Panama], watching a nest of the Black-throated Trogon [*Trogon rufus*], a Dusky-capped Flycatcher sunned itself in a patch of sunshine that fell on a prostrate tree close beside me. Lying flat on the trunk, it spread its wings and fluffed out its body feathers to let the sun’s rays penetrate deeply into them.”

The behaviors exhibited by Great Crested Flycatchers on the vinyl cover of my outdoor spa (i.e., stationary with splayed wings and tail against the vinyl surface, head cocked upward, mouth gaped) are all similar to those detailed in Hauser (1957). Considering the six possible functions for voluntary sunbathing (Kennedy 1969), two can be immediately eliminated in this situation. The observations occurred during the warmest part of the day, thereby eliminating sunrise thermoregulation. The flycatchers were not wet prior to sunbathing, thereby eliminating feather drying. Whether uropygial gland secretions or molting were involved, is unknown.

Ectoparasite control seems a viable hypothesis for my observations of sunbathing by Great Crested Flycatchers. Blem and Blem (1992) observed Tree Swallows (*Tachycineta bicolor*), Violet-green Swallows (*T. thalassina*), and Barn Swallows (*Hirundo rustica*) sunbathing on a shiny aluminum roof. They found the behavior to be consistent with either ectoparasite removal or soothing of skin areas involved in molting. Blem and Blem (1993) conducted experiments to see if sunbathing had an effect on ectoparasite load. One group of swallows was treated with a pesticide and a control group was not. They found that the control group sunbathed substantially more than the pesticide-treated group (23 vs. 2 visits, respectively) during the two-week expected efficacy of the pesticide treatment. During the subsequent two-week period, sunbathing visits increased for both groups, but remained higher for the control (49 vs. 29 visits). Blem and Blem surmised that ectoparasite control was the best explanation for this behavior. The actual mechanism for ectoparasite control by sunbathing was unknown, but various hypotheses mentioned were that high temperatures might kill, immobilize, or discourage ectoparasites; high temperatures might possibly concentrate ectoparasites for removal by preening; or sunlight itself may have a pesticidal effect (Blem and Blem 1993). Control of feather-degrading bacteria has also been mentioned as a reason for sunbathing (Gunderson 2008). Saranathan and Burt (2007) found that sunlight inhibited the growth of feather-degrading bacilli, a common inhabitant on avian plumage.

I speculate that there is a direct relationship between temperature and ectoparasite control, and that by sunbathing on the vinyl surface rather than the cooler wooden deck and rails, the Great Crested Flycatchers sought the hottest surface they could find/tolerate.

Although there is photo documentation of sunbathing by Great Crested Flycatchers on the Internet (Brown 2011, Tallman 2011, Roberts 2015, Zee 2015, Bacquie 2016), my observation may be only the second account in the published literature.

ACKNOWLEDGMENTS

I thank Eugene J. Scarpulla for his assistance locating several publications. I also thank two anonymous reviewers for their helpful comments.

LITERATURE CITED

- Bacque, J. 2016. A Great-crested Flycatcher sunbathing. janine-images. Available at: <http://www.janine-images.com/blog/index.php?itemid=413>. Accessed 30 January 2017.
- Blem, C.R., and L.B. Blem. 1992. Some observations of sunbathing in swallows. *Journal of Field Ornithology* 63(1):53–56.
- Blem, C.R., and L.B. Blem. 1993. Do swallows sunbathe to control ectoparasites? An experimental test. *The Condor* 95(3):728–730.
- Brown, D. 2011. Sunbathing (thanks Kenny) -Great-Crested Flycatcher (*Myiarchus crinitus*) - thanks Liam. David Brown Photography. Available at: <http://www.panoramio.com/photo/55503620>. Accessed 30 January 2017.
- Cardiff, S.W., and D.L. Dittmann. 2000. Brown-crested Flycatcher (*Myiarchus tyrannulus*), The Birds of North America (P.G. Rodewald, Editor). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America: <https://birdsna.org/Species-Account/bna/species/bncfly>. DOI: 10.2173/bna.496. Accessed 30 January 2017.
- Cardiff, S.W., and D.L. Dittmann. 2002. Ash-throated Flycatcher (*Myiarchus cinerascens*), The Birds of North America (P.G. Rodewald, Editor). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America: <https://birdsna.org/Species-Account/bna/species/astfly>. DOI: 10.2173/bna.664. Accessed 30 January 2017.
- Curry-Lindahl, K. 1970. Spread-wing postures in Pelecaniformes and Ciconiiformes. *The Auk* 87(2):371–372.
- Dunn, J.L., and J. Alderfer. 2006. *National Geographic Field Guide to the Birds of North America* (fifth edition). National Geographic Society, Washington, DC. 503 pp.
- Goodwin, D. 1967. Some possible functions of sun-bathing in birds. *British Birds* 60(9):363–364.
- Gunderson, A.R. 2008. Feather-degrading bacteria: A new frontier in avian and host–parasite research? *The Auk* 125(4):972–979.

- Hauser, D.C. 1957. Some observations on sun-bathing in birds. *The Wilson Bulletin* 69(1):78–90.
- Horsfall, J. 1984. Sunbathing: Is it for the birds? *New Scientist* 103(1420):28–31.
- Lanyon, W.E. 1958. The motivation of sunbathing in birds. *The Wilson Bulletin* 70(3):280.
- Kennedy, R.J. 1969. Sunbathing behaviour of birds. *British Birds* 62(7):249–258.
- Miller, K.E., and W.E. Lanyon. 2014. Great Crested Flycatcher (*Myiarchus crinitus*), The Birds of North America (P.G. Rodewald, Editor). Ithaca: Cornell Lab of Ornithology. Available at: <https://birdsna.org/Species-Account/bna/species/grcfly>. DOI: 10.2173/bna.300. Accessed 30 January 2017.
- Roberts, R. 2015. Sunbathing Bird Series: Great Crested Flycatcher. Available at: <https://plus.google.com/+RhondaRoberts/posts/B7ZH1yvFay3>. Accessed 30 January 2017.
- Saranathan, V., and E.H. Burt, Jr. 2007. Sunlight on feathers inhibits feather-degrading bacteria. *The Wilson Journal of Ornithology* 119(2):239–245. Available at: http://www.academia.edu/1459142/Sunlight_on_feathers_inhibits_feather-degrading_bacteria. Accessed 24 March 2017.
- Skutch, A.F. 1960. *Life Histories of Central American Birds II*. Pacific Coast Avifauna Number 34. Cooper Ornithological Society, Berkeley, CA. 593 pp. Available at: https://sora.unm.edu/sites/default/files/journals/pca/pca_034.pdf. Accessed 30 January 2017.
- Tallman, D. 2011. Great Crested Flycatcher: Anting or Sunning. Dan Tallman's Bird Blog. Available at: <http://dantallmansbirdblog.blogspot.com/2011/07/great-crested-flycatcher-anting.html>. Accessed 30 January 2017.
- Twit, R.C., and J.C. Twit. 2002. Dusky-capped Flycatcher (*Myiarchus tuberculifer*), The Birds of North America (P. G. Rodewald, Editor). Ithaca: Cornell Lab of Ornithology. Available at: <https://birdsna.org/Species-Account/bna/species/ducfly>. DOI: 10.2173/bna.631. Accessed 30 January 2017.
- Zee, M[ama]. 2015. Great Crested Flycatcher sunbathing. Toronto and Southern Ontario Birding. Available at: <http://outdoorontario.net/birds/phpBB/viewtopic.php?f=14&t=15751>. Accessed 30 January 2017.

Cedar Waxwings (*Bombycilla cedrorum*) Observed Feeding on Flying Insects

Jan G. Reese

*P.O. Box 298, St. Michaels, Maryland 21663
reesejan@ymail.com*

Abstract: The frugivorous Cedar Waxwing (*Bombycilla cedrorum*) subsists most of year on a diet of fruit nutritionally rich in sugars, but during spring, replaces fruit with a substantial amount of protein-rich plant pollen and insect prey. Studies report waxwing pursuit of flying insects in flycatcher-like sallies, but details of the behavior are lacking or poorly described. I report here waxwings observed aerial feeding on swarming ants (Hymenoptera: Formicidae). The waxwings used short, ungraceful sallies, generally without turns, and landed on nearby branches as soon as possible. Feeding on swarming insects during March–June suggests an opportunistic behavior to replace a dietary requirement during a portion of the year when sugar-rich fruit are not available; however, reasons for the dietary change may not be fully understood.

Keywords: aerial feeding, *Bombycilla cedrorum*, Cedar Waxwing, *Prenolepis imparis*, swarming Winter Ants

The Cedar Waxwing (*Bombycilla cedrorum*) is one of the most frugivorous birds in North America, with fruit comprising 70–84% of its diet (Stiles 1984, Witmer 1996). However, during March–June when fruit is scarce or absent, their diet includes substantial amounts of insects and flower parts for reasons not clearly understood (Beal 1893, Witmer 1996). Most individuals glean insects from vegetation foliage, but several studies also report aerial sallies to capture flying insects (Crouch 1936, Bartlett 1956, Harlow 1971, Pinkowski 1976, McPherson 1987, Witmer 1996, Witmer et al. 2014). These sallies are likened to those of flycatchers, but give little or no definitive descriptions of the behavior. I give here a more comprehensive description of the behavior, the prey, and potential reasons for the seasonal change in the diet.

OBSERVATION

Walking from within a willow oak-loblolly pine (*Quercus phellos* L.-*Pinus taeda* L.) association forest (Brush et al. 1980) in St. Michaels, Talbot County, Maryland on 8 March 2016 at 1430 hours, with an ambient temperature of 18.3° C (69.4° F), a flock of 28 Cedar Waxwings suddenly landed on top of a leafless

sweetgum (*Liquidambar styraciflua* L.) at the forest edge directly in front of me. I quickly halted, becoming motionless.

Upon landing, individual waxwings flew down and dispersed throughout branches of the sweetgum and the interior of an approximately 10 m (33 ft) tall adjacent southern magnolia (*Magnolia grandiflora* L.). Magnolia leaves covered only the outer portion of the branches, thus the tree's interior was open except for the bare branches emanating from the trunk. Waxwings quickly flitted from branch to branch of the gum, but mostly within the shaded interior of the magnolia. They appeared to be chasing some minute flying prey.

I eventually caught glimpses of sparkling clear-wing movement of tiny flying insects that were the object of the waxwings interest. After visual acclimation to the lighting situation, I realized there were hundreds or possibly thousands of tiny insects flitting about, perhaps staging or swarming within the magnolia branches and foliage.

The method used by waxwings to capture flying insects in the tree interior took several forms. It began from a branch posture of nearly 45° where waxwings appeared to be visually trying to locate aerial prey in front of them. Birds then simply darted forward, snatched the insects in the air, and continued nearly straight-ahead to another branch within the tree's interior. Others darted forward to snap prey in the air, and then turned abruptly to return and land near or adjacent to the departure branch. Still others, perched near openings in the exterior foliage, used a similar method, i.e., they darted outside the foliage to snatch an aerial insect then came back within the tree interior on openings lower in the foliage to land on a bare branch.

Ungraceful short sorties ranged up to approximately 3 m (10 ft) both inside and outside the tree foliage. Near midair collisions suggest limited abilities and/or unfamiliarity with this type of feeding behavior. Birds were silent during the 25-minute feeding event throughout the interior of the tree, with activity descending within 2 m (7 ft) of the ground and/or the observer.

After the feeding activity ceased and the waxwings departed, I netted 20 of the tiny (~3 mm [~0.1 in] long) flying insects for later identification. I forwarded the specimens for identification to Samuel E. Droege and Eugene J. Scarpulla at the Bee Inventory and Monitoring Laboratory (BIML) at the United States Geological Survey's Patuxent Wildlife Research Center in Beltsville, Maryland, where the insects were determined to be male alate (winged) ants, mounted, and photographed (Figure 1). The ant specimens were forwarded for species identification to Maryland ant specialist Timothy Foard at i2L Research USA, Inc. (i2L) in Baltimore, Maryland where they were identified as Winter Ants, *Prenolepis imparis* (Say) (Hymenoptera: Formicidae: Formicinae).




Figure 1. Winter Ant, *Prenolepis imparis* (Say) (Hymenoptera: Formicidae: Formicinae). Alate male. Body length: ~3.0 mm (mean, median, and mode; n = 13; measured to closest 0.5 mm, mouth hypognathous). **Top left:** frontal view; **top right:** lateral view; **bottom:** dorsal view. Specimens collected by Jan G. Reese in St. Michaels, Talbot County, Maryland on 8 March 2016 and photographed by Samuel W. Droege on 19 July and 9 September 2016.

I returned to the magnolia tree at the same time on three subsequent balmy days, but never again saw the insects or waxwings.

DISCUSSION

Observed aerial feeding in mid-afternoon on a late winter day was synonymous with the warmest period of the day when flying insects are most likely active. Aerial feeding has been reported in other studies that also took place during the warmest part of the day, 1100–1700 hours, from mid-March to September (Bartlett 1956, Baird and Meyerriecks 1965, Harlow 1971, Pinkowski 1976).

Waxwing aerial feeding has been observed most often in June–July (Bartlett 1956, Harlow 1971, Witmer 1996). Many of these events took place over ponds and streams which had emerging stoneflies (Plecoptera) or mayflies (Ephemeroptera) (Crouch 1936, Bartlett 1956, Pinkowski 1976, Witmer et al. 2014), or at other insect mating swarms and dispersal of flying ants (Baird and Meyerriecks 1965, Muller and Berger 1965, Harlow 1971). All of these occurrences suggest opportunistic feeding. Beal (1893) believed waxwings consume some quantity of insects at all times if they are easily obtained. Beal's observation suggested opportunistic aerial feeding since it was done in late winter, earlier than other studies, and a plentiful crop of American holly (*Ilex opaca* Aiton) was within view of the observed aerial feeding. Similarly, a study of waxwing winter fruit preference in Norman, Oklahoma in 1984–1985 found American holly berries of low preference when notable consumption of insects started in March (McPherson 1987).

Other waxwing studies also describe slower and less graceful movements than flycatchers or other species (Harlow 1971, Witmer et al. 2014), shorter flights (Pinkowski 1976) seldom returning to the origin branch (Bartlett 1956), and disregard for close proximity of the observer (Crouch 1936). Conversely, one study reports frequent returns to the origin branch (Harlow 1971). Waxwings have been credited with “hovering” to harvest fruit (Witmer et al. 2014) and “fluttering” while pursuing flying prey (Bartlett 1956). It appears both of these behaviors could be an adventitious strategy in aerial feeding. I observed no hovering or fluttering. On one occasion however, a waxwing appeared to miss the intended flying prey, thrust its wings into a vertical position in order to stop forward movement while simultaneously snapping or grabbing for the prey a second time. The bird then lost altitude and forward momentum, suddenly dropped, repositioned its wings, turned, and landed on a convenient branch. This behavior may be related to the fluttering observed by Bartlett (1956). The spirit, agility, and lack of finesse in Cedar Waxwing aerial feeding may be related to limb proportions, wing-loading ability, feather and bill structure, visibility, or other morphological features in comparison to flycatchers (Fitzpatrick 1985, Norberg 1986, Witmer 1996, Warrick 1998).

Winter Ants are commonly found throughout the continental United States. Nests are usually located in damp soils of habitats such as forests, forest edges, fallow fields, and around buildings (Klotz et al. 2008, Ellison et al. 2012). The species is cryophilic and commences foraging with temperatures around 0° C (32° F), thus it is one of the first ants to become active early in the year with nuptial flights commonly occurring by March–April (Klotz et al. 2008). Colonies actively forage in late-winter to mid-spring and mid-fall to early-winter while aestivating during the warmer months (Fisher and Cover 2007, Klotz et al. 2008).

Alate ants comprised seven percent of all insects found in Cedar Waxwing stomachs collected in 1885–1950 (Beal 1893, Witmer 1996). Many of those ants appeared to belong to the genus *Camponotus* Mayr (Formicidae: Formicinae). In Wisconsin during 1958–1963, prey in nine incidences of aerial-feeding waxwings during 28 August–22 September involved *Lasius alienus americanus* (Foerster) (Formicidae: Formicinae) (Muller and Berger 1965). Similarly, 13 species of birds, including Cedar Waxwings, were identified aerial-feeding on *L. alienus* on 23 August 1961 in Massachusetts (Baird and Meyerriecks 1965).

Waxwings appear to function well and maintain body mass for extended periods during the year on a ripe fruit diet nutritionally rich in various sugars; however during spring, consumption of fruit diminishes for several weeks while being replaced by a substantial amount of high protein plant pollen and insect prey (Witmer 1996, Witmer et al. 2014). A total of 212 waxwing stomachs collected in 1885–1950, mostly from eastern North America, contained insects in April through November being most frequent during May–June (Beal 1893, Witmer 1996). Flower parts (stamens and petals) comprised 44% of the waxwings diet in May and insects 41% when fruit crops were at a low or not available (Witmer 1996, Witmer et al. 2014). These observations further support the notion that a March–June slack period of fruit availability leads to supplemental feeding of plant and insect protein to replace a dietary requirement in preparation for migration, reproductive activities, or until preferred ripe fruit are again plentiful in July.

For instance, a hypothesis suggests in the temperate zone, adult frugivorous species such as waxwings have evolutionarily switched to a largely protein diet during the reproductive season lessening pressure on plants to produce fruit during the early and/or shorter growing season (Morton 1973). Additionally, frugivorous nestlings fed a largely protein diet during this time fledged in a shorter period of time than those on a fruit diet, while the shorter nestling period also reduces predation risk for nestlings.

Witmer (2001) noted nutritional sugary fruit of European cranberrybush (*Viburnum opulus* L.) in New York persisted through the winter, to be eaten

primarily by waxwings in spring when other fruits were not available. The waxwings also ate at this time nutritional, protein-rich, male catkins of eastern cottonwood (*Populus deltoides* W. Bartram ex Marshall). Further investigation found secondary compounds in the pulp of aged cranberrybush fruit made it acidic. The study suggested that in spring, waxwings consume foods rich in protein to produce bicarbonates as a buffer to the acid in the dehydrated fruit. Thus, hypotheses and waxwing studies indicate the spring dietary addition of protein-rich foods is still not fully understood.

In summary, numerous studies report March–September waxwing aerial feeding on swarming insects. The behavior is believed opportunistic and/or to meet a dietary requirement at a time of year when fruit are scarce or not available. In this observation, waxwing aerial feeding was generally a slow, straight, short, ungraceful sally followed by a landing at the nearest convenient branch rather than returning to the origin branch. The prey was from a mating swarm of alated Winter Ants.

ACKNOWLEDGMENTS

I thank Samuel W. Droege and Eugene J. Scarpulla (BIML) for mounting and photographing male alate ants and for comments on an earlier draft. Timothy Foard (i2L) generously identified the ant species. The manuscript benefited from constructive reviews by Elizabeth R. Clark and three anonymous reviewers.

LITERATURE CITED

- Baird, J., and A.J. Meyerriecks. 1965. Birds feeding on an ant mating swarm. *The Wilson Bulletin* 77(1):89–91.
- Bartlett, L.M. 1956. Observations on birds “hawking” insects. *The Auk* 73(1):127–128.
- Beal, F.E.L. 1893. Food habits of the Cedar Bird (*Ampelis cedrorum*). Pages 197–200 in: *Report of the Secretary of Agriculture - 1892*. Government Printing Office, Washington, DC. 656 pp.
- Brush, G.S., C. Lenk, and J. Smith. 1980. The natural forests of Maryland: an explanation of the vegetation map of Maryland. *Ecological Monographs* 50(1):77–92.
- Crouch, J.E. 1936. Nesting habits of the Cedar Waxwing (*Bombycilla cedrorum*). *The Auk* 53(1):1–8.

- Ellison, A.M., N.J. Gotelli, E.J. Farnsworth, and G.D. Alpert. 2012. *A Field Guide to the Ants of New England*. Yale University Press, New Haven, CT. 398 pp.
- Fisher, B.L., and S.P. Cover. 2007. *Ants of North America: A guide to the genera*. University of California Press, Berkeley and Los Angeles, CA. 194 pp.
- Fitzpatrick, J.W. 1985. Form, foraging behavior, and adaptive radiation in the Tyrannidae. *Ornithological Monographs* No. 36:447–470.
- Harlow, R.A., Jr. 1971. Birds feeding on an ant mating swarm in Maine. *The Wilson Bulletin* 83(2):201–202.
- Klotz, J., L. Hansen, R. Pospischil, and M. Rust. 2008. *Urban Ants of North America: Identification, biology, and management*. Cornell University Press, Ithaca, NY. 196 pp.
- McPherson, J.M. 1987. A field study of winter fruit preferences of Cedar Waxwings. *The Condor* 89(2):293–306.
- Morton, E.S. 1973. On the evolutionary advantages and disadvantages of fruit eating in tropical birds. *The American Naturalist* 107(953):8–22.
- Mueller, H.C., and D.D. Berger. 1965. Ring-billed Gulls feed on flying ants. *The Auk* 82(3):504.
- Norberg, U.M. 1986. Evolutionary convergence in foraging niche and flight morphology in insectivorous aerial-hawking birds and bats. *Ornis Scandinavica* 17(3):253–260.
- Pinkowski, B.C. 1976. Cedar Waxwings and Eastern Bluebirds feeding on winter stoneflies. *The Wilson Bulletin* 88(3):508–509.
- Stiles, E.W. 1984. Fruit for all seasons. *Natural History* 93(8):42–53.
- Warrick, D.R. 1998. The turning- and linear-maneuvering performance of birds: the cost of efficiency for coursing insectivores. *Canadian Journal of Zoology* 76(6):1063–1079.
- Witmer, M.C. 1996. Annual diet of Cedar Waxwings based on U.S. Biological Survey records (1885–1950) compared to diet of American Robins: contrasts in dietary patterns and natural history. *The Auk* 113(2):414–430.
- Witmer, M.C. 2001. Nutritional interactions and fruit removal: Cedar Waxwing consumption of *Viburnum opulus* fruits in spring. *Ecology* 82(11):3120–3130.

Witmer, M.C., D.J. Mountjoy, and L. Elliot. 2014. Cedar Waxwing (*Bombycilla cedrorum*). The Birds of North America Online (A. Poole, Editor) Cornell Lab of Ornithology, Ithaca, NY. Available at: <http://bna.birds.cornell.edu/bna/species/309>. DOI:10.2173/bna.309. [Updated URL: <https://birdsna.org/Species-Account/bna/species/cedwax>.]



“A Berry Good Day” by Ellen Lawler

Ellen painted this watercolor (the original is in color) of Cedar Waxwings in March 2017. For references, she used two photographs taken along the C & O Canal in Cumberland, Maryland, in October 2015.

2016 Maryland Mid-winter Bird Count: Corrigendum

Mary Anne Fluke

Nottingham, Maryland; mfluke62@gmail.com

The narrative of the 2016 Maryland Mid-winter Bird Count article (Fluke 2016), mistakenly reported a Boat-tailed Grackle for Howard County. The sighting should have been reported as a Baltimore Oriole.

Table 3 of the article also showed one Boat-tailed Grackle for Howard County that should have been recorded as one Baltimore Oriole. Hence, there were no Boat-tailed Grackles on the entire Mid-winter Bird Count, and alternatively there was one Baltimore Oriole. The corrected data lines for the blackbirds in Table 3 should read as follows:

| Species | C&O | FAAS | ACAS | FR | CA | HO | BA | HA | KE |
|----------------------|-----|------|------|-------|------|------|-----|-----|--------|
| Red-winged Blackbird | | 65 | | 1 155 | 80 | 1114 | 228 | 270 | 1032 |
| Eastern Meadowlark | | | | 14 | | 1 | 1 | 16 | |
| Rusty Blackbird | | | | | | | 4 | 5 | |
| Common Grackle | | | | | 4001 | 134 | 39 | 215 | 10,273 |
| Brown-headed Cowbird | | | | 60 | 4 | 211 | | 129 | 71 |
| Baltimore Oriole | | | | | | | 1 | | |
| blackbird sp. | | | | 300 | | | 8 | | |

LITERATURE CITED

Fluke, M.A. 2016. 2016 Maryland Mid-winter Bird Count. *Maryland Birdlife* 65(2):24–28.

2016 Maryland May Count

Marilyn E. Veek

8714 Reichs Ford Road, Frederick, Maryland 21704
mveek@yahoo.com

The annual Maryland May Count was held on Saturday, 14 May 2016. Results were submitted from 22 of Maryland’s 23 counties, two more than submitted results in 2015 (Table 1).

Weather statewide on 14 May was clear and calm in the morning, with a front accompanied by thunderstorms, gusty winds, and rain moving through from west to east in the afternoon. Temperatures were as low as 36° F (2° C) in Garrett County in the morning, to as high as 77° F (25° C) in Dorchester County at 4:11 p.m. just before the front arrived. Calvert and Dorchester reported wind gusts up to 45–50 mph (72–80 kph) with the front.

Table 1. 2016 Maryland May Count: participating counties, county codes, and compilers. (The dashed line separates the Western and Eastern Shore counts.)

| County | Code | Compiler |
|-----------------|------|---------------------------------|
| Garrett | GA | Nadine Jakubowski |
| Allegany | AL | J.B. Churchill |
| Washington | WA | Mark Abdy |
| Frederick | FR | David Smith |
| Montgomery | MO | Diane Ford |
| Carroll | CA | Don Jewell |
| Howard | HO | Kevin Heffernan & Chuck Stirrat |
| Baltimore | BA | Joel Martin |
| Prince George’s | PG | Fred Fallon |
| Anne Arundel | AA | Dotty Mumford |
| Charles | CH | Lynne Wheeler |
| Calvert | CT | Sherman Suter |
| St. Mary’s | SM | J. Tyler Bell |
| Harford | HA | Rick Cheicante |
| Cecil | CE | Richard Donham |
| Kent | KE | Walter Ellison |
| Talbot | TA | David Palmer |
| Caroline | CN | Debby Bennett |
| Dorchester | DO | Harry Armistead |
| Somerset | SO | Paul Bystrak |
| Wicomico | WI | Travis Clemens |
| Worcester | WO | Marcia Balestri |

A total of 461 counters in 279 parties reported 244 species and 127,262 individual birds (Tables 2 and 3).

Among the most notable sightings reported were: 115 Ruddy Turnstones (9 in Dorchester, 4 in Kent, 102 in Worcester), 9 Purple Sandpipers (Worcester), 218 Northern Gannett (Anne Arundel), 213 Brown Pelicans (1 in Calvert, 52 in St. Mary's, 140 in Dorchester, and 20 in Wicomico), 1 White Ibis (Worcester), 1 Swallow-tailed Kite (Howard) (Figure 1), 1 Loggerhead Shrike (Garrett) (report submitted to MD/DC Records Committee), 34 Gray-cheeked Thrushes (ranging from 1 to 8 each in 12 different counties), 322 Swainson's Thrushes (a nice rebound from only 22 individuals last year; reported from 17 counties with Howard high at 106), 4604 Cedar Waxwings (more than 3 times higher than previous years, reported from every county except Somerset, with a Howard high at 1049), 121 Canada Warblers (up from a low of 7 last year; reported from 18 counties with Howard high at 29), 17 Dickcissels (9 in Allegany, 1 in Carroll, 3 in Prince George's (Figure 2), 1 in Harford, 3 in Kent) (Tables 4 and 5).

Additionally, 2 Swainson's Warblers were reported from Calvert. The report was submitted to the MD/DC Records Committee, but was not accepted by the Committee.

Notable misses this year were Blue-winged Teal, Northern Shoveler, and Black-necked Stilt. White-crowned Sparrows continue to decrease, with only 4 individuals reported (1 from Garrett, 2 from Allegany, 1 from Howard).

The 2015 May Count was notable for low numbers of most thrushes and eighteen species of warblers. This year showed improved numbers, especially for Gray-cheeked and Swainson's Thrushes, as noted above. Counts for all eighteen species of warblers noted as low in 2015 were higher, as well. The cases in which 2016 numbers were lower than 2015 were not large differences, and in two cases were decreases from counts noted as significantly high in 2015: Prothonotary Warbler (123 in 2016 vs. 184 in 2015) and Yellow-throated Warbler (92 vs. 153.)

Comparing data from 2003, 2013, 2014, 2015, and 2016 there have been clear increases in Black and Turkey Vultures, Common Raven, Bald Eagle, and Brown-headed Nuthatch. Blue Jay numbers this year were about twice the numbers reported in previous years. Raptor numbers were improved this year compared to 2015, for example, 502 Bald Eagles vs. 401.

This was a good year in terms of the increase in number of individual birds seen, counties reporting, and counters involved in the May Count, but a few large population counties would do much better with more people in the field.

Table 2. 2016 Maryland May Count: Western Shore summary.

| | GA | AL | WA | FR | MO | CA | HO | BA | PG | AA | CH | CT | SM | HA |
|-------------------|-------|------|------|-------|------|-------|--------|-------|------|------|------|-------|-------|------|
| Parties | 24 | 6 | 10 | 12 | 1 | 12 | 34 | 42 | 28 | 21 | 8 | 10 | 15 | 10 |
| Observers | 42 | 6 | 18 | 21 | 4 | 18 | 70 | 44 | 49 | 30 | 13 | 16 | 15 | 33 |
| Start time | 0530 | 0540 | 0430 | 0445 | 0700 | 0500 | 0445 | 0000 | 0530 | 0530 | 0530 | 0000 | 0300 | 0005 |
| End time | 0000 | 1900 | 1900 | 1930 | 1400 | 2100 | 2330 | 2156 | 2030 | 1930 | 2000 | 2018 | 2030 | 2045 |
| Driving | | | | | | | | | | | | | | |
| hours | 61.15 | 24 | 32 | 47.25 | | 27 | 34.8 | 1 | 53.3 | 7.5 | 21 | 4.5 | 85.82 | 35.5 |
| miles | 464 | 112 | 283 | 481 | | 268.5 | 426.4 | 24 | 108 | 49.5 | 268 | 23.75 | 324.2 | 169 |
| Walking | | | | | | | | | | | | | | |
| hours | 60.15 | 18 | 24.5 | 35.5 | | 43 | 190.1 | 73.5 | 91 | 68.5 | 37.5 | 49.45 | 18.75 | 43.5 |
| miles | 48.75 | 17 | 18.5 | 23.2 | 5.5 | 29.1 | 143.5 | 72.76 | 59 | 61 | 31 | 29.75 | 16.5 | 21.5 |
| Owling | | | | | | | | | | | | | | |
| hours | 3.75 | | 1.5 | 1.25 | | 0.75 | 1.8 | | 4 | | 1 | 1.67 | 2.25 | |
| miles | 43 | | 3 | | | 0 | 5 | | 5 | | 0.5 | 6 | 64 | |
| Stationary | | | | | | | | | | | | | | |
| hours | 23 | | 5.5 | 6.5 | 0.25 | 18 | 4 | 19.2 | 14 | 15.5 | 5 | 5.42 | 6.5 | |
| Other conveyance | | | | | | | | | | golf | | | | |
| tractor | | | | | | | | | | cart | bike | | | |
| hours | 3.55 | | | | | | | | | 2 | 3 | | | |
| miles | 20.5 | | | | | | | | | 3 | 3 | | | |
| Total species | 147 | 136 | 121 | 144 | 77 | 130 | 148 | 139 | 141 | 147 | 123 | 147 | 138 | 151 |
| Total individuals | 7445 | 2725 | 4237 | 8842 | 452 | 5904 | 14,732 | 4624 | 6672 | 9597 | 5476 | 6090 | 6232 | 6493 |

Table 3. 2016 Maryland May Count: Eastern Shore and total summary.

| | CE | KE | TA | CN | DO | SO | WI | WO | West & East Total |
|-------------------|------|------|------|-----------|--------|-------|------|-------|----------------------|
| Parties | 4 | 5 | 10 | 11 | 9 | 2 | 5 | | 279 |
| Observers | 6 | 7 | 14 | 16 | 14 | 2 | 8 | 15 | 461 |
| Start time | 0438 | 0520 | 0400 | 0430 | 0000 | 0500 | 0508 | 0500 | |
| End time | 1500 | 2045 | 2000 | 2130 | 2100 | 1730 | 1530 | 1952 | |
| Driving | | | | | | | | | |
| hours | | 18.5 | 42 | 30 | 69 | 4.5 | 4.35 | 30.15 | 628.97 |
| miles | | 141 | 108 | 292 | 584 | 8.1 | 26.3 | 350 | 4510.75 |
| Walking | | | | | | | | | |
| hours | 16.5 | 15 | 12 | 29.5 | 20 | 2 | 7 | 20.35 | 875.8 |
| miles | 13 | 12.1 | 6.5 | 13 | 16 | 2.5 | 3.5 | 20.6 | 664.26 |
| Owling | | | | | | | | | |
| hours | | | 2 | 2.75 | 8 | | | 0.5 | 31.22 |
| miles | | | 27 | 44 | 32 | | | 2 | 231.5 |
| Stationary | | | | | | | | | |
| hours | 0.5 | | 5.25 | 13 | 1.5 | 3.583 | 6 | 3.7 | 156.403 |
| Other conveyance | | | | golf cart | boat | | | | |
| hours | | | | 1 | 2 | | | | 11.55 |
| miles | | | | 1 | | | | | 27.5 |
| Total species | 100 | 135 | 133 | 131 | 166 | 90 | 95 | 161 | 244 |
| Total individuals | 1467 | 5798 | 3607 | 7760 | 10,215 | 1603 | 1323 | 5968 | 127,262 |



Figure 1. Swallow-tailed Kite (*Elanoides forficatus*). Alpha Ridge Landfill, Howard County, Maryland, 14 May 2016, photographed by Joe Hanfman.



Figure 2. Dickcissel (*Spiza americana*). Beltsville Agricultural Research Center, Prince George's County, Maryland, 14 May 2016, photographed by Jack Saba.

Table 4. 2016 Maryland May Count: Western Shore observed species.

(* = report submitted to MD/DC Records Committee)

| Species | GA | AL | WA | FR | MO | CA | HO | BA | PG | AA | CH | CT | SM | HA |
|---------------------------|-----|-----|-----|-----|----|-----|-----|-----|-----|-----|-----|----|-----|-----|
| Canada Goose | 316 | 86 | 182 | 329 | 43 | 331 | 674 | 116 | 259 | 267 | 181 | 71 | 112 | 284 |
| Mute Swan | 65 | | | 1 | | | | | | | | | 10 | 9 |
| Tundra Swan | | | | 1 | | | | | | | | | | 1 |
| Wood Duck | | 28 | 9 | 24 | 1 | 2 | 40 | 5 | 95 | 43 | 32 | 14 | 2 | 23 |
| Gadwall | | | | | | | | | | | | | | |
| American Black Duck | | | | | 2 | | | | | | 2 | | 1 | 2 |
| Mallard | 63 | 27 | 85 | 95 | 4 | 31 | 82 | 64 | 57 | 74 | 28 | 57 | 40 | 25 |
| Green-winged Teal | | | | | | | | | | | | | | |
| Canvasback | | | | | | | | | | | | | | |
| Ring-necked Duck | | | | 1 | | | | | | | | | | |
| Lesser Scaup | 1 | | | | | | | | | 3 | | | | |
| unidentified scaup | | | | | | | | | 1 | | | | | |
| Surf Scoter | | | | | | | | | | | | | | |
| Long-tailed Duck | | | | | | | | | | | | | | |
| Bufflehead | 2 | | | 2 | | 1 | | | | | | | | |
| Hooded Merganser | | | | 10 | | | | | | 4 | | | | 3 |
| Common Merganser | 5 | | | 1 | 1 | | | | | | | | | |
| Red-breasted Merganser | | | | | | | | | | | 1 | 1 | | |
| Ruddy Duck | | | 2 | | | | | | | | | 8 | 21 | |
| unidentified duck | | | | | | | | | | | | | | |
| Northern Bobwhite | | | | 1 | | | | | | | | | | |
| Ruffed Grouse | 7 | 1 | | | | | | | | | | | | |
| Wild Turkey | 39 | 11 | 19 | 2 | | 5 | 12 | | 11 | 4 | 20 | 10 | 8 | 32 |
| Pied-billed Grebe | | 2 | | | | | 5 | 1 | | | | | 1 | |
| Horned Grebe | | | | | | | | | | | | | | |
| Rock Pigeon | 58 | 132 | 88 | 87 | 1 | 41 | 86 | 44 | 22 | 36 | 1 | | 16 | 33 |
| Mourning Dove | 151 | 36 | 111 | 195 | 12 | 107 | 263 | 85 | 117 | 139 | 114 | 71 | 68 | 124 |
| Yellow-billed Cuckoo | 3 | 7 | 15 | 19 | 1 | 9 | 17 | 7 | 26 | 12 | 2 | 8 | 6 | 21 |
| Black-billed Cuckoo | 3 | 2 | | 1 | | 1 | 2 | | | | | | | 1 |
| Common Nighthawk | | | | | | | 10 | 26 | 4 | 19 | | 4 | | 7 |
| Chuck-will's-widow | | | | | | | | | | | | 3 | 6 | |
| Eastern Whip-poor-will | 3 | | 5 | | | | | 1 | | | | | 1 | 20 |
| Chimney Swift | 23 | 39 | 45 | 111 | 13 | 61 | 127 | 104 | 95 | 118 | 83 | 43 | 20 | 81 |
| Ruby-throated Hummingbird | 60 | 8 | 7 | 13 | 2 | 16 | 15 | 17 | 8 | 14 | 13 | 10 | 18 | 14 |
| Clapper Rail | | | | | | | | | | | | 1 | 1 | |
| King Rail | | | | | | | | | | | | | | 2 |
| Virginia Rail | | | | | | 2 | | | | | | 2 | | 3 |
| Sora | | | | | | | | | | 1 | | 1 | 1 | 4 |
| unidentified rail | | | | | | | | | | | | | | |
| Common Gallinule | | 1 | | 1 | | | | 1 | | | | | | |
| American Coot | | 3 | 1 | 2 | | | 1 | 2 | 1 | | | | | |
| Sandhill Crane | 4 | | | | | | | | | | | | | |
| American Oystercatcher | | | | | | | | | | | | | | |
| Black-bellied Plover | | | | | | | | | | | | | | |
| Semipalmated Plover | 7 | | | | | | | | | | | 3 | 18 | 1 |
| unidentified plover | | | | | | | | | | | | | | |
| (mostly Black-bellied?) | | | | | | | | | | | | | | |
| Killdeer | 34 | 12 | 16 | 32 | | 8 | 10 | 1 | 8 | 4 | 16 | 6 | 14 | 42 |
| Ruddy Turnstone | | | | | | | | | | | | | | |

| Species | GA | AL | WA | FR | MO | CA | HO | BA | PG | AA | CH | CT | SM | HA |
|--|----|----|----|-----|----|----|-----|----|-----|-----|-----|-----|-----|-----|
| Red Knot | | | | | | | | | | | | | | |
| Sanderling | | | | | | | | | | | | | | 1 |
| Dunlin | | | | | | | | | | | | | | |
| Purple Sandpiper | | | | | | | | | | | | | | |
| Least Sandpiper | 3 | 9 | | | | | 4 | 15 | 1 | 2 | | 24 | 69 | 21 |
| White-rumped Sandpiper | 7 | | | | | | | | | | | | | |
| Pectoral Sandpiper | | | | | | | | | | | | | | |
| Semipalmated Sandpiper | | 1 | | | | 2 | | | | | | 4 | 10 | 6 |
| unidentified sandpiper (mostly Dunlin?) | | | | | | | | | | | | | | |
| Short-billed Dowitcher | | | | | | | | | | | | 1 | | |
| Wilson's Snipe | 1 | | | | | | 7 | | 1 | | | | | |
| American Woodcock | 4 | | 8 | | | | 2 | | | | | | | 7 |
| Spotted Sandpiper | 27 | 14 | 13 | 15 | 2 | 8 | 37 | 16 | 26 | 22 | 12 | 22 | 11 | 13 |
| Solitary Sandpiper | 9 | 2 | 1 | 10 | 1 | 5 | 33 | 4 | 15 | 5 | 1 | 4 | 9 | 21 |
| Greater Yellowlegs | | | | 1 | | | | | 5 | | | 2 | 1 | 1 |
| Willet | | | | | | | | | | | | | | |
| Lesser Yellowlegs | 1 | 4 | 1 | | | | | | 2 | | 1 | | | 1 |
| Bonaparte's Gull | | | 3 | | | | | | | | | 14 | 50 | |
| Laughing Gull | | | | | | | | | 16 | 8 | 3 | 31 | 50 | |
| Ring-billed Gull | | | | | | 3 | | 18 | 2 | 61 | 176 | 162 | 527 | 20 |
| Herring Gull | | | | | | | | 6 | | 49 | 4 | 41 | 54 | |
| Great Black-backed Gull | | | | | | | | 3 | | 27 | | 75 | 20 | |
| unidentified gull. | | | | | | | | 5 | 1 | | | | 15 | |
| Least Tern | | | | | | | | 1 | | 10 | | 2 | | 14 |
| Caspian Tern | | | | | | | | 1 | 2 | 3 | 2 | | | 5 |
| Common Tern | | | | | | | | 2 | | 4 | 26 | 1 | | |
| Forster's Tern | | | | | | | | | | | 4 | | | |
| Royal Tern | | | | | | | | | | | | | | |
| unidentified tern | | | | | | | | | 2 | | | | 5 | |
| Black Skimmer | | | | | | | | | | | | | | |
| Common Loon | 5 | 2 | 3 | | | 10 | 2 | 2 | 1 | 3 | 4 | | 1 | |
| Northern Gannet | | | | | | | | | | 218 | | | | |
| Double-crested Cormorant | 17 | | 26 | 3 | 1 | 5 | 63 | 29 | 160 | | 163 | 371 | 181 | 492 |
| American White Pelican | | | | | | | | | | | | | | |
| Brown Pelican | | | | | | | | | | | | 1 | 52 | |
| American Bittern | | | | | | | 2 | | | | | | | |
| Least Bittern | | | | | | | | | | 1 | | | 1 | 3 |
| Great Blue Heron | 1 | 2 | 8 | 20 | 9 | 20 | 37 | 20 | 62 | 72 | 39 | 20 | 34 | 72 |
| Great Egret | | | | | | | 1 | 1 | 3 | 2 | 1 | | 4 | 2 |
| Snowy Egret | | | | | | | | 1 | | 2 | | 10 | | |
| Little Blue Heron | | | | | | | | 3 | | | | | | |
| Tricolored Heron | | | | | | | | | | | | | | |
| Cattle Egret | | | | | | | | | | 4 | | | | |
| Green Heron | 8 | 5 | 1 | 8 | 2 | 4 | 26 | 4 | 8 | 8 | 5 | 3 | 4 | 6 |
| Black-crowned Night-Heron | | | | 11 | | | 1 | | | | | 1 | | |
| Yellow-crowned Night-Heron | | | | | | 2 | | 4 | | | | | | |
| White Ibis | | | | | | | | | | | | | | |
| Glossy Ibis | | | | | | | | | | | | 5 | | 1 |
| Black Vulture | | 2 | 16 | 33 | | 86 | 106 | 8 | 56 | 35 | 119 | 19 | 55 | 28 |
| Turkey Vulture | 69 | 43 | 61 | 103 | 2 | 86 | 225 | 46 | 77 | 199 | 75 | 90 | 58 | 136 |
| Osprey | 1 | 1 | 4 | 5 | | 2 | 26 | 15 | 53 | 101 | 176 | 77 | 119 | 51 |
| Swallow-tailed Kite | | | | | | | 1 | | | | | | | |

| Species | GA | AL | WA | FR | MO | CA | HO | BA | PG | AA | CH | CT | SM | HA |
|-------------------------------|-----|----|-----|-----|----|-----|-----|-----|-----|------|-----|-----|-----|-----|
| Mississippi Kite | | | | | | | | | | | | | | 1 |
| Bald Eagle | 5 | 1 | | 2 | | 5 | 13 | 5 | 36 | 62 | 53 | 7 | 25 | 51 |
| Northern Harrier | | | 1 | 1 | | | 2 | 2 | 3 | 6 | | 1 | | 3 |
| Sharp-shinned Hawk | | | | | | 1 | 2 | 2 | 2 | 49 | 2 | 1 | | 2 |
| Cooper's Hawk | 1 | 1 | 1 | 1 | | 8 | 6 | 3 | 4 | 9 | | 3 | 1 | 2 |
| Red-shouldered Hawk | | | 6 | 15 | 1 | 14 | 72 | 21 | 24 | 15 | 9 | 9 | 7 | 12 |
| Broad-winged Hawk | 6 | | 1 | 3 | 1 | | 8 | 1 | 2 | 136 | 1 | | 2 | 4 |
| Red-tailed Hawk | 8 | 2 | 12 | 23 | | 18 | 19 | 9 | 21 | 15 | 8 | 8 | 2 | 18 |
| unidentified hawk | | | | | | | | | | | | | | |
| Barn Owl | | | | | | | | | | | 16 | | | |
| Eastern Screech-Owl | 1 | 3 | | 4 | | 1 | 1 | | 1 | 2 | 1 | 5 | | 1 |
| Great Horned Owl | 1 | | | | | 1 | 6 | | 1 | 1 | | | | 1 |
| Barred Owl | 5 | 1 | 2 | 5 | | 4 | 13 | 1 | 4 | 6 | 3 | 13 | 1 | 7 |
| Belted Kingfisher | 1 | 2 | 3 | 5 | | 6 | 15 | 7 | 5 | 7 | 4 | 5 | 2 | 4 |
| Red-headed Woodpecker | 13 | | | 6 | | 6 | 3 | 3 | 1 | | 7 | 12 | 9 | 5 |
| Red-bellied Woodpecker | 38 | 22 | 43 | 135 | 9 | 84 | 237 | 63 | 69 | 70 | 56 | 62 | 28 | 49 |
| Yellow-bellied Sapsucker | 6 | | | | | | | | | | | | | |
| Downy Woodpecker | 36 | 12 | 27 | 45 | 11 | 44 | 98 | 35 | 31 | 37 | 11 | 28 | 15 | 23 |
| Hairy Woodpecker | 33 | 5 | 7 | 5 | 3 | 10 | 19 | 14 | 9 | 7 | 4 | 11 | 3 | 6 |
| Northern Flicker | 39 | 12 | 8 | 13 | 1 | 17 | 31 | 9 | 18 | 10 | 4 | 8 | 10 | 18 |
| Pileated Woodpecker | 19 | 10 | 20 | 23 | 1 | 16 | 49 | 10 | 16 | 13 | 11 | 18 | 11 | 14 |
| American Kestrel | 6 | 1 | 3 | 8 | | 5 | 3 | | 5 | | 1 | | | 2 |
| Merlin | | | | | | | | | | 2 | | | | |
| Peregrine Falcon | | | | | | | | | | 1 | | | | |
| Olive-sided Flycatcher | | 2 | | | | | | | | | | | | |
| Eastern Wood-Pewee | 14 | 9 | 19 | 92 | 4 | 40 | 135 | 38 | 46 | 49 | 52 | 79 | 55 | 28 |
| Acadian Flycatcher | 2 | 5 | 11 | 51 | 1 | 36 | 116 | 6 | 44 | 42 | 53 | 88 | 50 | 31 |
| Alder Flycatcher | 1 | 1 | | | | | 1 | | 1 | | | | | |
| Willow Flycatcher | 1 | 1 | 1 | 5 | | 8 | 17 | 3 | | 1 | | | | 4 |
| Alder/Willow Flycatcher | | | | | | | | | | | | | | |
| (Traill's Flycatcher) | | | | | | | | | | | | | | |
| Least Flycatcher | 16 | 2 | | 1 | | | 2 | 3 | 1 | | | 1 | | |
| unidentified <i>Empidonax</i> | | | | 4 | | 2 | | | | | | | | |
| Eastern Phoebe | 25 | 10 | 19 | 36 | | 32 | 61 | 23 | 19 | 6 | 14 | 10 | 10 | 23 |
| Great Crested Flycatcher | 13 | 8 | 43 | 91 | 5 | 41 | 93 | 25 | 49 | 45 | 49 | 54 | 68 | 45 |
| Eastern Kingbird | 17 | 10 | 20 | 67 | 1 | 46 | 103 | 30 | 104 | 77 | 38 | 60 | 37 | 70 |
| Loggerhead Shrike* | 1 | | | | | | | | | | | | | |
| White-eyed Vireo | 2 | 4 | 4 | 13 | | 24 | 50 | 11 | 43 | 23 | 42 | 58 | 46 | 35 |
| Yellow-throated Vireo | 1 | 2 | 6 | 9 | 1 | 6 | 14 | 15 | 24 | 4 | 14 | 16 | 3 | 23 |
| Blue-headed Vireo | 33 | 2 | | 3 | 1 | 2 | 2 | 4 | 1 | 9 | | | 2 | |
| Philadelphia Vireo | | | | | | | | 2 | | | | | | |
| Warbling Vireo | | 9 | 14 | 33 | 4 | 16 | 14 | 31 | 3 | 2 | | | | 16 |
| Red-eyed Vireo | 236 | 43 | 86 | 235 | 26 | 113 | 388 | 109 | 154 | 195 | 122 | 190 | 211 | 94 |
| Blue Jay | 178 | 78 | 114 | 159 | 7 | 154 | 562 | 292 | 110 | 2244 | 44 | 48 | 59 | 115 |
| American Crow | 277 | 75 | 147 | 159 | 3 | 186 | 383 | 86 | 70 | 107 | 107 | 97 | 158 | 108 |
| Fish Crow | | 4 | 17 | 36 | 1 | 14 | 53 | 16 | 29 | 37 | 18 | 27 | 43 | 14 |
| unidentified crow | | | | 30 | 1 | | 137 | 9 | | 5 | | 7 | 23 | 8 |
| Common Raven | 26 | 11 | 2 | 4 | | 2 | 2 | 5 | 1 | 1 | | | | 1 |
| Horned Lark | | | 11 | 9 | | 1 | 16 | 1 | | | | 8 | 10 | 2 |
| Purple Martin | 34 | | 32 | 28 | | 22 | 83 | 3 | 28 | 16 | 128 | 30 | 82 | 10 |
| Tree Swallow | 298 | 25 | 42 | 71 | 2 | 74 | 125 | 53 | 196 | 168 | 40 | 32 | 15 | 181 |
| N. Rough-winged Swallow | 85 | 15 | 11 | 35 | 3 | 17 | 32 | 32 | 13 | 22 | 10 | 13 | 2 | 46 |
| Bank Swallow | 4 | 1 | | 1 | | 1 | | 1 | 4 | 11 | | | | 6 |

| Species | GA | AL | WA | FR | MO | CA | HO | BA | PG | AA | CH | CT | SM | HA |
|-------------------------------------|-----|-----|-----|-----|----|-----|------|-----|-----|-----|-----|-----|-----|-----|
| Cliff Swallow | 149 | 70 | 46 | 16 | | 51 | 39 | | 1 | 2 | | | | |
| Barn Swallow | 620 | 25 | 96 | 162 | 4 | 156 | 248 | 58 | 107 | 208 | 168 | 89 | 163 | 248 |
| Carolina Chickadee | | | 64 | 144 | 9 | 83 | 261 | 64 | 83 | 65 | 50 | 119 | 104 | 78 |
| Black-capped Chickadee | 117 | 22 | 9 | | | | | | | | | | | |
| Tufted Titmouse | 68 | 36 | 80 | 119 | 6 | 97 | 260 | 73 | 83 | 84 | 82 | 114 | 97 | 54 |
| Red-breasted Nuthatch | 6 | 5 | | | | | | | | | | | | |
| White-breasted Nuthatch | 39 | 17 | 28 | 41 | 5 | 30 | 72 | 46 | 26 | 23 | 12 | 22 | 9 | 17 |
| Brown-headed Nuthatch | | | | | | | | | | | | 4 | 13 | |
| Brown Creeper | 9 | 3 | | 1 | | | | | | | | | | |
| House Wren | 56 | 20 | 23 | 100 | 5 | 53 | 99 | 37 | 12 | 18 | 2 | 3 | 6 | 20 |
| Winter Wren | 1 | 1 | | | | | | | | | | | | |
| Sedge Wren | 1 | | | | | | 1 | | | | | | | |
| Marsh Wren | 1 | | | | | | | 1 | | 4 | 4 | 3 | 4 | 3 |
| Carolina Wren | 9 | 34 | 56 | 157 | 14 | 54 | 165 | 57 | 79 | 60 | 67 | 86 | 101 | 44 |
| Blue-gray Gnatcatcher | 15 | 11 | 35 | 95 | 3 | 78 | 192 | 46 | 109 | 95 | 64 | 138 | 38 | 84 |
| Golden-crowned Kinglet | 4 | 1 | | | | | | | | | | | | |
| Ruby-crowned Kinglet | 3 | | | 1 | | | 12 | 3 | | 2 | | 4 | | |
| Eastern Bluebird | 31 | 64 | 94 | 95 | | 86 | 180 | 29 | 79 | 62 | 36 | 36 | 42 | 80 |
| Veery | 13 | 1 | 1 | 1 | | 15 | 19 | 11 | 3 | 2 | | 10 | 6 | 7 |
| Gray-cheeked Thrush | | | | 2 | | | 8 | 2 | 3 | | 1 | 6 | 3 | 2 |
| Swainson's Thrush | 1 | | 6 | 13 | 1 | 7 | 106 | 49 | 22 | 22 | 5 | 28 | 7 | 29 |
| Hermit Thrush | 11 | 1 | | | | 1 | 6 | | | | | | | |
| unidentified <i>Catharus</i> thrush | | | | | | | | | | | | | | |
| Wood Thrush | 24 | 13 | 56 | 137 | 2 | 103 | 244 | 51 | 65 | 55 | 53 | 73 | 54 | 25 |
| American Robin | 428 | 207 | 176 | 569 | 19 | 186 | 520 | 227 | 270 | 328 | 162 | 107 | 148 | 171 |
| Gray Catbird | 138 | 82 | 127 | 277 | 19 | 301 | 438 | 149 | 87 | 16 | 17 | 78 | 98 | 177 |
| Brown Thrasher | 37 | 11 | 36 | 35 | | 31 | 26 | 5 | 23 | 12 | 45 | 20 | 34 | 13 |
| Northern Mockingbird | | 17 | 58 | 97 | 1 | 57 | 155 | 26 | 55 | 70 | 90 | 48 | 75 | 68 |
| European Starling | 265 | 180 | 393 | 801 | 7 | 307 | 594 | 208 | 413 | 207 | 306 | 83 | 222 | 440 |
| Cedar Waxwing | 21 | 24 | 28 | 210 | 42 | 244 | 1049 | 480 | 427 | 942 | 136 | 431 | 88 | 189 |
| House Sparrow | 78 | 65 | 109 | 187 | 26 | 103 | 236 | 94 | 66 | 50 | 140 | 78 | 72 | 59 |
| American Pipit | | | | | | | | | 1 | | | | | |
| House Finch | 40 | 37 | 47 | 95 | 5 | 100 | 144 | 54 | 41 | 52 | 59 | 55 | 28 | 40 |
| Purple Finch | 25 | 3 | 3 | | | | | | | | | | | |
| Pine Siskin | 11 | 3 | | | | | | | | | | | | |
| American Goldfinch | 213 | 84 | 104 | 226 | 7 | 227 | 461 | 97 | 111 | 277 | 64 | 94 | 124 | 93 |
| Evening Grosbeak | | 3 | | | | | | | | | | | | |
| Ovenbird | 122 | 13 | 28 | 91 | 1 | 58 | 158 | 25 | 71 | 52 | 42 | 52 | 48 | 35 |
| Worm-eating Warbler | 1 | 10 | 5 | 9 | | 5 | 9 | 1 | 3 | 4 | 2 | 21 | 9 | 3 |
| Louisiana Waterthrush | 9 | 2 | 4 | 15 | 1 | 5 | 27 | 4 | 4 | 5 | 16 | 13 | 6 | 11 |
| Northern Waterthrush | 3 | 10 | | 2 | | 1 | 9 | 7 | 7 | 5 | 2 | 7 | 2 | 11 |
| Golden-winged Warbler | 2 | | | 1 | | | | | | | | | | |
| Blue-winged Warbler | | 1 | 1 | 1 | | 11 | 10 | | 1 | 3 | | | | 3 |
| Black-and-white Warbler | 37 | 3 | 9 | 24 | 2 | 7 | 34 | 19 | 11 | 17 | 8 | 35 | 26 | 12 |
| Prothonotary Warbler | 1 | | 2 | 5 | | | | | 6 | 12 | 8 | 1 | | 5 |
| Tennessee Warbler | 1 | 3 | | 1 | | | 12 | 1 | | 1 | | | | |
| Nashville Warbler | | | | | | | 3 | 3 | | | | 1 | | |
| Mourning Warbler | 1 | 1 | 1 | | | 1 | | 1 | 1 | | | | | |
| Kentucky Warbler | 5 | 2 | 1 | 4 | | 2 | 12 | 4 | | 1 | 11 | 8 | 10 | 10 |
| Common Yellowthroat | 196 | 26 | 27 | 76 | 2 | 104 | 265 | 60 | 128 | 84 | 80 | 147 | 128 | 99 |
| Hooded Warbler | 10 | 14 | 5 | 2 | | 2 | 8 | 4 | 5 | 6 | 29 | 47 | 12 | 3 |
| American Redstart | 68 | 29 | 26 | 46 | | 41 | 151 | 67 | 65 | 25 | 22 | 61 | 36 | 67 |
| Cape May Warbler | | | | 4 | | 1 | 8 | 3 | 3 | 1 | | 2 | | 4 |

| Species | GA | AL | WA | FR | MO | CA | HO | BA | PG | AA | CH | CT | SM | HA |
|------------------------------|-----|-----|-----|-----|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Cerulean Warbler | 7 | 8 | 15 | 1 | | | 3 | | | | | 2 | | 4 |
| Northern Parula | 19 | 4 | 5 | 21 | 4 | 18 | 137 | 43 | 48 | 39 | 76 | 114 | 41 | 44 |
| Magnolia Warbler | 35 | 5 | 5 | 5 | 1 | 1 | 26 | 28 | 24 | 16 | 2 | 30 | 26 | 28 |
| Bay-breasted Warbler | | 3 | | 6 | | 2 | 6 | 1 | 5 | | 8 | | 5 | 2 |
| Blackburnian Warbler | 24 | | | 2 | | | 5 | 9 | 3 | | 5 | | 4 | 2 |
| Yellow Warbler | 71 | 35 | 7 | 34 | 2 | 60 | 72 | 33 | 16 | 18 | 11 | 14 | 13 | 55 |
| Chestnut-sided Warbler | 101 | 4 | 3 | 3 | | 2 | 23 | 23 | 13 | | 9 | | 8 | 3 |
| Blackpoll Warbler | 1 | 10 | 11 | 24 | 5 | 9 | 38 | 19 | 74 | 28 | 19 | 80 | 34 | 29 |
| Black-throated Blue Warbler | 36 | 3 | 1 | 11 | | | 18 | 76 | 37 | 16 | 11 | 1 | 53 | 13 |
| Palm Warbler | | 1 | | | 2 | 1 | 2 | | 1 | 3 | | | | 1 |
| Pine Warbler | 1 | 7 | 4 | 2 | | 1 | 2 | 4 | 5 | 3 | 5 | 16 | 59 | 2 |
| Yellow-rumped Warbler | 9 | 2 | 15 | 35 | 2 | 16 | 77 | 56 | 38 | 22 | 6 | 17 | 2 | 44 |
| Yellow-throated Warbler | | | 2 | 1 | | | 2 | | 2 | 3 | 8 | 33 | 8 | 1 |
| Prairie Warbler | | 5 | 16 | 6 | | 13 | 24 | 2 | 14 | 12 | 11 | 8 | 28 | 23 |
| Black-throated Green Warbler | 54 | 6 | 3 | 9 | | 2 | 16 | 11 | 11 | 9 | 2 | 25 | 2 | 5 |
| Canada Warbler | 8 | | 2 | 1 | 1 | 4 | 29 | 14 | 15 | 12 | 4 | 7 | 1 | 10 |
| Wilson's Warbler | 2 | | | 1 | | | 1 | 3 | 2 | 1 | 2 | 5 | 3 | 2 |
| Yellow-breasted Chat | 1 | | 4 | 6 | | 7 | 12 | 4 | 9 | 10 | 4 | 8 | 7 | 30 |
| unidentified warbler | | | | | | | | | | | | | | |
| Eastern Towhee | 162 | 36 | 33 | 83 | 1 | 61 | 149 | 13 | 47 | 31 | 15 | 39 | 45 | 26 |
| Chipping Sparrow | 146 | 30 | 84 | 160 | 3 | 125 | 231 | 34 | 110 | 66 | 106 | 104 | 131 | 97 |
| Field Sparrow | 52 | 4 | 39 | 70 | | 59 | 42 | 5 | 18 | 4 | 20 | 7 | 12 | 12 |
| Vesper Sparrow | 1 | 2 | | 6 | | | 2 | | | | | | | |
| Savannah Sparrow | 7 | 3 | | 2 | | 6 | 20 | | 4 | 9 | | 7 | 3 | 8 |
| Grasshopper Sparrow | 10 | 2 | 9 | 11 | | 9 | 11 | | 12 | 3 | 9 | 11 | 48 | 18 |
| Henslow's Sparrow | | 2 | | | | | | | | | | | | |
| Saltmarsh Sparrow | | | | | | | | | | | | | | |
| Seaside Sparrow | | | | | | | | 3 | | | | 1 | 2 | |
| Song Sparrow | 189 | 67 | 48 | 134 | 5 | 106 | 123 | 22 | 35 | 65 | 7 | 40 | 13 | 25 |
| Lincoln's Sparrow | | 2 | | | | | 1 | | 2 | 1 | | 2 | 21 | |
| Swamp Sparrow | 40 | 1 | 1 | | | 1 | 2 | 4 | 2 | 1 | 6 | 7 | 3 | 18 |
| White-throated Sparrow | 2 | 4 | 7 | 10 | 1 | 26 | 41 | 37 | 4 | 11 | 2 | 10 | 5 | 15 |
| White-crowned Sparrow | 1 | 2 | | | | | 1 | | | | | | | |
| unidentified sparrow | | | | | | | | | | | | | | |
| Dark-eyed Junco | 6 | | | | | | | | | | | | | |
| Summer Tanager | | | | | | | | | 4 | 4 | 8 | 42 | 31 | |
| Scarlet Tanager | 84 | 12 | 16 | 64 | 4 | 47 | 137 | 34 | 64 | 23 | 53 | 33 | 30 | 61 |
| Northern Cardinal | 92 | 103 | 119 | 471 | 21 | 271 | 659 | 162 | 219 | 256 | 169 | 198 | 237 | 186 |
| Rose-breasted Grosbeak | 63 | 10 | 5 | 5 | | 3 | 13 | 9 | 1 | 6 | 1 | 7 | 1 | 3 |
| Blue Grosbeak | | | | 7 | | | 4 | 2 | 22 | 10 | 11 | 18 | 20 | 24 |
| Indigo Bunting | 73 | 39 | 86 | 191 | 4 | 63 | 228 | 38 | 119 | 102 | 147 | 108 | 127 | 102 |
| Dickcissel | | 9 | | | | 1 | | | 3 | | | | | 1 |
| Bobolink | 72 | 33 | 12 | 34 | | 3 | 180 | | 87 | 105 | | 31 | 19 | 75 |
| Red-winged Blackbird | 400 | 83 | 55 | 407 | 8 | 183 | 618 | 70 | 482 | 436 | 270 | 256 | 237 | 244 |
| Eastern Meadowlark | 40 | 28 | 15 | 33 | | 11 | 12 | | 9 | 2 | 5 | 13 | 35 | 18 |
| Rusty Blackbird | 1 | | | | | | | | | | | | | |
| Common Grackle | 117 | 79 | 228 | 361 | 7 | 76 | 231 | 87 | 155 | 176 | 328 | 205 | 431 | 117 |
| Boat-tailed Grackle | | | | | | | | | | | | | | |
| Brown-headed Cowbird | 44 | 28 | 63 | 69 | 5 | 72 | 152 | 34 | 46 | 96 | 116 | 60 | 29 | 95 |
| unidentified blackbird | | | | | | | | | 13 | | | | | |
| Orchard Oriole | 16 | 11 | 5 | 15 | 3 | 18 | 53 | 16 | 39 | 16 | 38 | 28 | 13 | 67 |
| Baltimore Oriole | 89 | 19 | 72 | 76 | 8 | 76 | 127 | 40 | 19 | 7 | 5 | 6 | 1 | 38 |

Table 5. 2016 Maryland May Count: Eastern Shore and total observed species. (* = report submitted to MD/DC Records Committee)

| Species | CE | KE | TA | CN | DO | SO | WI | WO | Total |
|---|----|-----|-----|-----|-----|-----|----|-----|-------|
| Canada Goose | 14 | 81 | 79 | 177 | 305 | | 46 | 239 | 4192 |
| Mute Swan | | | | | | | | | 85 |
| Tundra Swan | | | | | 1 | | | | 3 |
| Wood Duck | | 23 | 7 | 27 | 22 | 1 | 2 | 1 | 401 |
| Gadwall | | | | 2 | | | | 1 | 3 |
| American Black Duck | | 2 | 2 | 2 | 27 | 5 | | 19 | 64 |
| Mallard | | 104 | 83 | 38 | 168 | 3 | 5 | 37 | 1170 |
| Green-winged Teal | | | 2 | | 1 | | | | 3 |
| Canvasback | | 1 | | | 3 | | | | 4 |
| Ring-necked Duck | | | | | | | | | 1 |
| Lesser Scaup | | | | | | | | | 4 |
| unidentified scaup | | | | | | | | 1 | 2 |
| Surf Scoter | | | 3 | | | | | | 3 |
| Long-tailed Duck | | | 9 | | | | | | 9 |
| Bufflehead | | | | | | | | 5 | 10 |
| Hooded Merganser | | 3 | | | | | | | 20 |
| Common Merganser | | | | | | | | | 7 |
| Red-breasted Merganser | | | | | 6 | | | | 8 |
| Ruddy Duck | | 3 | | | 47 | | | | 81 |
| unidentified duck | | | | | 2 | | | | 2 |
| Northern Bobwhite | | | | 5 | 74 | | | | 80 |
| Ruffed Grouse | | | | | | | | | 8 |
| Wild Turkey | 1 | 13 | 20 | 66 | 23 | 1 | 5 | 5 | 307 |
| Pied-billed Grebe | | | | | | | | | 9 |
| Horned Grebe | | | 2 | | | | | | 2 |
| Rock Pigeon | 10 | 9 | 10 | 36 | 34 | | 1 | 15 | 760 |
| Mourning Dove | 24 | 65 | 39 | 128 | 93 | 5 | 25 | 20 | 1992 |
| Yellow-billed Cuckoo | 7 | 10 | 3 | 13 | 30 | 2 | 17 | 25 | 260 |
| Black-billed Cuckoo | | | | | | | | | 10 |
| Common Nighthawk | | 1 | | | 1 | | | 2 | 74 |
| Chuck-will's-widow | | | 2 | 1 | 21 | | 3 | 7 | 43 |
| Eastern Whip-poor-will | 8 | | | 3 | | | 1 | 2 | 44 |
| Chimney Swift | 4 | 82 | 49 | 402 | 71 | | 1 | 21 | 1593 |
| Ruby-throated Hummingbird | | 10 | 3 | 18 | 14 | 1 | 13 | 9 | 283 |
| Clapper Rail | | | | | 10 | 4 | | 2 | 18 |
| King Rail | | | 2 | | 3 | | | | 7 |
| Virginia Rail | | 3 | 3 | 1 | 36 | 2 | | 3 | 55 |
| Sora | | | | | | | | 1 | 8 |
| unidentified rail | | | | | 4 | | | | 4 |
| Common Gallinule | | | | | 12 | | | | 15 |
| American Coot | | | | | | | | | 10 |
| Sandhill Crane | | | | | | | | | 4 |
| American Oystercatcher | | | | | 2 | | | 19 | 21 |
| Black-bellied Plover | | 221 | | 50 | 2 | 18 | | 638 | 929 |
| Semipalmated Plover | 4 | 14 | 275 | 3 | 11 | | | 9 | 345 |
| unidentified plover (mostly Black-bellied?) | | | | | | 150 | | | 150 |
| Killdeer | 2 | 4 | 8 | 29 | 23 | | 3 | 6 | 278 |
| Ruddy Turnstone | | 4 | | | 9 | | | 102 | 115 |
| Red Knot | | | | | | | | 6 | 6 |

| Species | CE | KE | TA | CN | DO | SO | WI | WO | Total |
|---|-----|-----|-----|-----|-----|-----|----|-----|-------|
| Sanderling | | | | | 4 | | | 3 | 8 |
| Dunlin | | | 79 | | 270 | 263 | | 756 | 1368 |
| Purple Sandpiper | | | | | | | | 9 | 9 |
| Least Sandpiper | 2 | 36 | 66 | 59 | 124 | 15 | 3 | 89 | 542 |
| White-rumped Sandpiper | | | | | 1 | | | | 8 |
| Pectoral Sandpiper | | | 3 | | | 11 | | 1 | 15 |
| Semipalmated Sandpiper | 1 | 3 | | | 11 | 47 | | 27 | 112 |
| unidentified sandpiper (mostly Dunlin?) | | | | | | 450 | | | 450 |
| Short-billed Dowitcher | | | | | | 46 | | 52 | 99 |
| Wilson's Snipe | | | | 10 | | | | | 19 |
| American Woodcock | | | | 2 | 1 | | | | 24 |
| Spotted Sandpiper | 2 | 5 | 8 | 12 | 24 | | 2 | 11 | 302 |
| Solitary Sandpiper | | 1 | 1 | 18 | 2 | | | 3 | 145 |
| Greater Yellowlegs | | 3 | 1 | 12 | 18 | 20 | 1 | 12 | 77 |
| Willet | | | 1 | | 18 | 14 | | 30 | 63 |
| Lesser Yellowlegs | | 6 | 1 | 5 | 22 | 3 | | 44 | 91 |
| Bonaparte's Gull | | 5 | 10 | 3 | 1 | | | | 86 |
| Laughing Gull | | 1 | 33 | 661 | 443 | 60 | 27 | 505 | 1838 |
| Ring-billed Gull | 2 | 279 | 15 | 410 | 20 | | | 20 | 1715 |
| Herring Gull | | | 51 | 3 | 75 | 7 | | 128 | 418 |
| Great Black-backed Gull | | 1 | | 2 | 17 | 2 | | 81 | 228 |
| unidentified gull | | | | | 4 | | | | 25 |
| Least Tern | | 4 | 8 | | 6 | 1 | | 15 | 61 |
| Caspian Tern | 1 | 7 | | | | | | | 21 |
| Common Tern | | 2 | 6 | 2 | 4 | 3 | | 80 | 130 |
| Forster's Tern | | | 33 | 1 | 19 | 8 | 39 | 30 | 134 |
| Royal Tern | | | | | 4 | 13 | | 125 | 142 |
| unidentified tern | | | | | 47 | | | | 54 |
| Black Skimmer | | | | | | | | 9 | 9 |
| Common Loon | | | | | 4 | | | 2 | 39 |
| Northern Gannet | | | | | | | | | 218 |
| Double-crested Cormorant | 242 | 32 | 514 | 38 | 598 | 14 | 92 | 373 | 3414 |
| American White Pelican | | | | | 1 | | | | 1 |
| Brown Pelican | | | | | 140 | | 20 | | 213 |
| American Bittern | | | | 1 | | | | | 3 |
| Least Bittern | | | | | 2 | | | | 7 |
| Great Blue Heron | 41 | 32 | 54 | 43 | 105 | 2 | 7 | 3 | 703 |
| Great Egret | | 1 | 9 | | 59 | 4 | 1 | 25 | 113 |
| Snowy Egret | | | 5 | 1 | 29 | 16 | | 19 | 83 |
| Little Blue Heron | | | 1 | | | | | 3 | 7 |
| Tricolored Heron | | | | | 1 | | | | 1 |
| Cattle Egret | | | 9 | 4 | | | | 22 | 39 |
| Green Heron | | 5 | 7 | 8 | 3 | | | 4 | 119 |
| Black-crowned Night-Heron | | | | | 3 | | | 2 | 18 |
| Yellow-crowned Night-Heron | | | | | | | | 1 | 7 |
| White Ibis | | | | | | | | 1 | 1 |
| Glossy Ibis | | 4 | 19 | 1 | 1 | | | 59 | 90 |
| Black Vulture | 25 | 39 | 21 | 40 | 51 | | 12 | 14 | 765 |
| Turkey Vulture | 18 | 152 | 85 | 184 | 247 | 18 | 52 | 85 | 2111 |
| Osprey | | 1 | 116 | 103 | 43 | 171 | 28 | 6 | 1099 |
| Swallow-tailed Kite | | | | | | | | | 1 |
| Mississippi Kite | 1 | | | | | | | | 2 |
| Bald Eagle | 9 | 29 | 18 | 26 | 138 | 3 | 9 | 5 | 502 |

| Species | CE | KE | TA | CN | DO | SO | WI | WO | Total |
|--|----|-----|-----|-----|-----|----|----|-----|-------|
| Northern Harrier | 2 | | | | | | | | 21 |
| Sharp-shinned Hawk | | 1 | | | 1 | | | | 63 |
| Cooper's Hawk | | 1 | 3 | 3 | 3 | 1 | | | 51 |
| Red-shouldered Hawk | 4 | 11 | | 3 | 3 | 1 | 1 | 4 | 232 |
| Broad-winged Hawk | 2 | 1 | | | 1 | | | | 169 |
| Red-tailed Hawk | 6 | 10 | 4 | 12 | 9 | 1 | | 1 | 206 |
| unidentified hawk | | | | | 1 | | | | 1 |
| Barn Owl | | | | | 1 | | | | 17 |
| Eastern Screech-Owl | | | 2 | 3 | 4 | | 2 | | 31 |
| Great Horned Owl | | | 2 | 6 | 8 | | | | 27 |
| Barred Owl | | 9 | 5 | 7 | | | 2 | 1 | 89 |
| Belted Kingfisher | 1 | | 2 | 2 | 4 | | 1 | | 76 |
| Red-headed Woodpecker | | | | | 18 | | 1 | 4 | 88 |
| Red-bellied Woodpecker | 36 | 51 | 15 | 55 | 18 | 1 | 23 | 18 | 1182 |
| Yellow-bellied Sapsucker | | | | | | | | | 6 |
| Downy Woodpecker | 2 | 18 | 8 | 17 | 24 | 2 | 7 | 6 | 537 |
| Hairy Woodpecker | 1 | 9 | 4 | 4 | 4 | 2 | | 1 | 161 |
| Northern Flicker | 3 | 4 | 3 | 7 | 12 | 3 | 1 | 1 | 232 |
| Pileated Woodpecker | 5 | 11 | 8 | 24 | 6 | 2 | 4 | 4 | 295 |
| American Kestrel | | | | 4 | | | | | 38 |
| Merlin | | | | | | | | 1 | 3 |
| Peregrine Falcon | | | | | | | | | 1 |
| Olive-sided Flycatcher | | | | | | | | | 2 |
| Eastern Wood-Pewee | 17 | 48 | 8 | 37 | 32 | 5 | 11 | 1 | 819 |
| Acadian Flycatcher | 11 | 50 | 4 | 8 | 8 | 1 | 12 | 21 | 651 |
| Alder Flycatcher | | | | | | | | | 4 |
| Willow Flycatcher | | | 1 | | | | | | 42 |
| Alder/Willow Flycatcher (Traill's Flycatcher) | | | | | | | | 1 | 1 |
| Least Flycatcher | | 1 | | | | | | | 27 |
| unidentified <i>Empidonax</i> | | | | | | | | | 6 |
| Eastern Phoebe | 4 | 5 | 3 | 23 | 2 | | 2 | 3 | 330 |
| Great Crested Flycatcher | 18 | 82 | 43 | 87 | 145 | 10 | 25 | 46 | 1085 |
| Eastern Kingbird | 2 | 28 | 23 | 26 | 79 | 2 | 14 | 19 | 873 |
| Loggerhead Shrike* | | | | | | | | | 1 |
| White-eyed Vireo | 13 | 20 | 4 | 17 | 40 | 8 | 21 | 45 | 523 |
| Yellow-throated Vireo | 3 | 2 | | | 1 | 1 | | 9 | 154 |
| Blue-headed Vireo | | | | | | | | | 59 |
| Philadelphia Vireo | | | | | | | | | 2 |
| Warbling Vireo | | 6 | | 6 | | | | | 154 |
| Red-eyed Vireo | 68 | 126 | 19 | 53 | 97 | 2 | 11 | 47 | 2625 |
| Blue Jay | 25 | 45 | 17 | 79 | 23 | | 10 | 15 | 4378 |
| American Crow | 43 | 30 | 62 | 105 | 134 | 6 | 40 | 51 | 2434 |
| Fish Crow | 7 | 20 | 15 | 23 | 28 | 3 | 4 | 29 | 438 |
| unidentified crow | | | | | 12 | | | | 232 |
| Common Raven | | | | | | | | | 55 |
| Horned Lark | | 15 | 12 | 63 | 7 | | | 19 | 174 |
| Purple Martin | | 120 | 111 | 167 | 432 | 3 | 41 | 100 | 1470 |
| Tree Swallow | 20 | 50 | 46 | 41 | 251 | 8 | 3 | 34 | 1775 |
| Northern Rough-winged Swallow | 2 | 6 | 3 | 8 | 5 | | | | 360 |
| Bank Swallow | | 1 | 1 | | 21 | | | | 52 |
| Cliff Swallow | | | 3 | 6 | 4 | | | | 387 |
| Barn Swallow | 4 | 96 | 125 | 175 | 275 | 11 | 31 | 67 | 3136 |

| Species | CE | KE | TA | CN | DO | SO | WI | WO | Total |
|-------------------------------------|----|-----|-----|-----|-----|----|----|-----|-------|
| Carolina Chickadee | 35 | 63 | 34 | 70 | 80 | 8 | 28 | 20 | 1462 |
| Black-capped Chickadee | | | | | | | | | 148 |
| Tufted Titmouse | 27 | 78 | 30 | 84 | 86 | 3 | 28 | 28 | 1617 |
| Red-breasted Nuthatch | | | | | | | | | 11 |
| White-breasted Nuthatch | 6 | 6 | 4 | 10 | | | | 1 | 414 |
| Brown-headed Nuthatch | | 3 | 5 | | 43 | 8 | 2 | 14 | 92 |
| Brown Creeper | | | | | 1 | | | | 14 |
| House Wren | 2 | 11 | 3 | 3 | 39 | 14 | 2 | 5 | 533 |
| Winter Wren | | | | | | | | | 2 |
| Sedge Wren | | | | | | | | | 2 |
| Marsh Wren | | 3 | 2 | 2 | 62 | 16 | 10 | | 115 |
| Carolina Wren | 9 | 83 | 19 | 49 | 90 | 10 | 47 | 30 | 1320 |
| Blue-gray Gnatcatcher | 9 | 60 | 4 | 13 | 67 | 11 | 19 | 55 | 1241 |
| Golden-crowned Kinglet | | | | | | | | | 5 |
| Ruby-crowned Kinglet | | | | | 1 | | | 1 | 27 |
| Eastern Bluebird | 2 | 23 | 21 | 71 | 28 | 1 | | 11 | 1071 |
| Veery | 8 | 2 | 1 | | 1 | | | | 101 |
| Gray-cheeked Thrush | 2 | 3 | | 1 | | | | 1 | 34 |
| Swainson's Thrush | 16 | 7 | | 2 | | | 1 | | 322 |
| Hermit Thrush | | | | 4 | | | | | 23 |
| unidentified <i>Catharus</i> thrush | | | | | 2 | | | | 2 |
| Wood Thrush | 71 | 64 | 21 | 31 | 13 | 1 | 11 | 21 | 1188 |
| American Robin | 13 | 298 | 113 | 384 | 181 | | 24 | 66 | 4597 |
| Gray Catbird | 45 | 64 | 25 | 48 | 51 | 9 | 9 | 67 | 2322 |
| Brown Thrasher | 5 | 21 | 20 | 30 | 14 | | 3 | 7 | 428 |
| Northern Mockingbird | 8 | 58 | 30 | 73 | 85 | 1 | 14 | 26 | 1112 |
| European Starling | 21 | 381 | 95 | 889 | 404 | 11 | 13 | 122 | 6362 |
| Cedar Waxwing | 13 | 55 | 26 | 29 | 110 | | 6 | 54 | 4604 |
| House Sparrow | 3 | 182 | 32 | 330 | 74 | | 1 | 3 | 1988 |
| American Pipit | | | | | | | | | 1 |
| House Finch | | 27 | 24 | 93 | 17 | | 9 | 9 | 976 |
| Purple Finch | | | | | | | | | 31 |
| Pine Siskin | | | | | | | | | 14 |
| American Goldfinch | 13 | 107 | 25 | 81 | 76 | 4 | 16 | 20 | 2524 |
| Evening Grosbeak | | | | | | | | | 3 |
| Ovenbird | 74 | 66 | 7 | 27 | 62 | 11 | 29 | 45 | 1117 |
| Worm-eating Warbler | 6 | 5 | 3 | 2 | 15 | 1 | 6 | 8 | 128 |
| Louisiana Waterthrush | | | 1 | 1 | 2 | | 3 | 6 | 135 |
| Northern Waterthrush | 1 | 8 | | | 2 | | | 5 | 82 |
| Golden-winged Warbler | | | | | | | | | 3 |
| Blue-winged Warbler | | | | | 1 | | | | 32 |
| Black-and-white Warbler | 3 | 20 | 9 | 5 | 26 | | 4 | 20 | 331 |
| Prothonotary Warbler | 1 | 4 | 2 | 15 | 20 | 2 | 8 | 31 | 123 |
| Tennessee Warbler | | | | | | | | | 19 |
| Nashville Warbler | | | | | | | | | 7 |
| Mourning Warbler | | 1 | | | | | | | 7 |
| Kentucky Warbler | | 9 | 2 | 2 | | | | 10 | 93 |
| Common Yellowthroat | 25 | 92 | 41 | 38 | 286 | 23 | 31 | 52 | 2010 |
| Hooded Warbler | 15 | | | | | | | 1 | 163 |
| American Redstart | 9 | 26 | 14 | 16 | 49 | | 4 | 34 | 856 |
| Cape May Warbler | | | | | 1 | | | 1 | 28 |
| Cerulean Warbler | | | | | | | | | 40 |
| Northern Parula | 5 | 26 | 9 | 10 | 20 | | 5 | 21 | 709 |

| Species | CE | KE | TA | CN | DO | SO | WI | WO | Total |
|------------------------------|----|-----|-----|-----|-----|----|-----|-----|-------|
| Magnolia Warbler | 3 | 7 | 6 | | 14 | | 2 | 11 | 275 |
| Bay-breasted Warbler | 2 | 1 | | | 3 | | | 1 | 45 |
| Blackburnian Warbler | | 2 | 1 | | 1 | | | 4 | 62 |
| Yellow Warbler | 2 | 4 | 7 | 3 | 13 | 1 | 1 | 12 | 484 |
| Chestnut-sided Warbler | | 6 | 1 | 1 | 2 | | | 2 | 211 |
| Blackpoll Warbler | 2 | 14 | 14 | 17 | 12 | 1 | 4 | 10 | 455 |
| Black-throated Blue Warbler | 3 | 5 | 5 | 6 | 9 | 2 | | 3 | 344 |
| Palm Warbler | | | | 3 | | | | | 14 |
| Pine Warbler | 8 | 9 | 16 | 3 | 167 | 17 | 12 | 20 | 363 |
| Yellow-rumped Warbler | 7 | 6 | 3 | 7 | 7 | | 2 | 4 | 377 |
| Yellow-throated Warbler | | 2 | | | 5 | 1 | 4 | 20 | 92 |
| Prairie Warbler | 2 | | 3 | 7 | 13 | 3 | 8 | 23 | 221 |
| Black-throated Green Warbler | 5 | 5 | | 1 | | 1 | | 2 | 169 |
| Canada Warbler | 3 | 6 | 2 | | 1 | | | 1 | 121 |
| Wilson's Warbler | | | | | 2 | | | | 24 |
| Yellow-breasted Chat | 3 | 6 | | 15 | 7 | 2 | 2 | 9 | 146 |
| unidentified warbler | | | | | 6 | | | | 6 |
| Eastern Towhee | 31 | 21 | 1 | 12 | 25 | 8 | 8 | 16 | 863 |
| Chipping Sparrow | 8 | 118 | 36 | 98 | 166 | 6 | 29 | 30 | 1918 |
| Field Sparrow | 13 | 5 | 5 | 15 | 23 | | 5 | 9 | 419 |
| Vesper Sparrow | | 2 | | 7 | | | | | 20 |
| Savannah Sparrow | | | 2 | | 2 | | | | 73 |
| Grasshopper Sparrow | | 9 | 10 | 15 | 43 | | 2 | 4 | 236 |
| Henslow's Sparrow | | | | | | | | | 2 |
| Saltmarsh Sparrow | | | | | 1 | | | 1 | 2 |
| Seaside Sparrow | | | | 112 | 43 | | | 22 | 183 |
| Song Sparrow | 7 | 12 | 1 | 9 | 11 | 5 | 1 | 5 | 930 |
| Lincoln's Sparrow | | | | | | | | | 29 |
| Swamp Sparrow | | | | | 12 | | | 1 | 99 |
| White-throated Sparrow | 3 | 4 | 1 | 4 | | | | 2 | 189 |
| White-crowned Sparrow | | | | | | | | | 4 |
| unidentified sparrow | | | | | 12 | | | | 12 |
| Dark-eyed Junco | | | | | | | | | 6 |
| Summer Tanager | | | 2 | 9 | 33 | 1 | 5 | 17 | 156 |
| Scarlet Tanager | 27 | 40 | 6 | 15 | 15 | | 2 | 14 | 781 |
| Northern Cardinal | 35 | 255 | 79 | 168 | 133 | 13 | 33 | 46 | 3925 |
| Rose-breasted Grosbeak | 1 | | 7 | 1 | 2 | | | 5 | 143 |
| Blue Grosbeak | | 24 | 20 | 49 | 74 | 4 | 9 | 16 | 314 |
| Indigo Bunting | 16 | 151 | 19 | 114 | 109 | 7 | 26 | 63 | 1932 |
| Dickcissel | | 3 | | | | | | | 17 |
| Bobolink | 30 | 1 | 3 | | 11 | | | 3 | 699 |
| Red-winged Blackbird | 41 | 398 | 212 | 320 | 921 | 46 | 104 | 155 | 5946 |
| Eastern Meadowlark | 5 | 2 | | 4 | 21 | 3 | | 8 | 264 |
| Rusty Blackbird | | | | | | | | | 1 |
| Common Grackle | 8 | 392 | 174 | 648 | 564 | 3 | 30 | 138 | 4555 |
| Boat-tailed Grackle | | | | | 13 | 7 | | 16 | 36 |
| Brown-headed Cowbird | 19 | 100 | 23 | 103 | 118 | 7 | 13 | 43 | 1335 |
| unidentified blackbird | | | | | 2 | | | | 15 |
| Orchard Oriole | 3 | 27 | 7 | 17 | 61 | 3 | 6 | 14 | 476 |
| Baltimore Oriole | 15 | 12 | 1 | 17 | 2 | | | 26 | 656 |

2016 Maryland Fall Count

Charles R. Stirrat

13318 Hunt Ridge, Ellicott City, Maryland 21042-1155

Stirrbird@outlook.com

The annual Maryland Fall Count is held on the third weekend in September with the choice of Saturday or Sunday at the discretion of the county and/or compiler. In 2016, eleven of Maryland's 23 counties submitted results. Five counties counted on 17 September, six on 18 September (Table 1). Receiving results from eleven counties is a decrease in participation for the Fall Count. For several counties this seasonal count has been a long held tradition, with this year being the 26th count for Allegany, 23rd for Howard, and 21st for Dorchester.

Table 1: 2016 Maryland Fall Count: participating counties, county codes, survey dates, and compilers.

| County | Code | Survey Date | Compiler |
|-----------------|------|-------------------|-----------------------------------|
| Allegany | AL | 18 September 2016 | Chuck Hager |
| Carroll | CA | 17 September 2016 | Don Jewell |
| Howard | HO | 17 September 2016 | Mike McClure, Chuck Stirrat |
| Baltimore | BA | 17 September 2016 | Joel Martin |
| Harford | HA | 18 September 2016 | Rick Cheicante |
| Cecil | CE | 18 September 2016 | Patricia Valdata, Sean McCandless |
| Prince George's | PG | 18 September 2016 | Dave Mozurkewich |
| Calvert | CT | 18 September 2016 | Sherman Suter |
| Kent | KE | 18 September 2016 | Walter Ellison |
| Caroline | CN | 17 September 2016 | Debby Bennett |
| Dorchester | DO | 17 September 2016 | Harry Armistead |

As would be expected weather varied over the extent of the state and from Saturday to Sunday. Temperatures were generally warmer on Sunday with highs in the mid 80s° F (~29° C), with the exception of Allegany at 79° F (26.1° C); whereas the highs on Saturday were around 80° F (26.7° C), with the exception of Dorchester at 86° F (30° C). Lows ranged from 55° F (12.8° C) in Caroline to 71° F (21.7° C) in Calvert. All reported calm to light winds, generally increasing as the day proceeded. Most reported partly cloudy conditions throughout the day. There were no reports of precipitation for either day, except for Allegany which had heavy rain on Sunday afternoon. Many compilers described the birding as slow with limited migrants.

A total of 190 field observers in 127 parties turned up 198 species and 65,447 individual birds. They spent a total of 473.6 hours and covered 334.2 mi [537.8 km] on foot and spent 26.0 hours stationary. Birders spent 135.9 hours and 1723.4 mi [2773.5 km] birding by car. Observers covered 10.5 mi [16.9 km] in 9.5 hours by other modes of transport, including by boat. Participants in 7 counties reported spending 18.7 hours traveling 23.7 mi [38.1 km] while owling. Feeder watchers spent 30.8 hours (Table 2).

Table 2. 2016 Maryland Fall Count: summary.

| | AL | CA | HO | BA | HA | CE | PG | CT | KT | CN | DO | Total |
|-------------------|-------|-------|-------|------|------|------|------|------|-------|-------|-------|--------|
| Total Species | 92 | 104 | 125 | 93 | 54 | 50 | 114 | 123 | 108 | 97 | 148 | 198 |
| Total Birds | 1572 | 6854 | 12651 | 1984 | 540 | 228 | 8340 | 7068 | 4173 | 6882 | 15185 | 65,447 |
| Start Time | 0500 | 0630 | 0615 | 0645 | 1130 | 0807 | 0530 | 0600 | 0530 | 0430 | 0430 | 0345 |
| Stop Time | 1830 | 1900 | 2000 | 1846 | 1745 | 1148 | 2300 | 1830 | 1930 | 2130 | 1930 | 23:45 |
| Parties | 8 | 11 | 47 | 16 | 3 | 1 | 11 | 8 | 3 | 9 | 10 | 127 |
| Individual People | 9 | 25 | 62 | 17 | 3 | 1 | 16 | 12 | 5 | 14 | 26 | 190 |
| Hours Driving | 10.0 | 18.5 | 25.8 | | | | 7.0 | 6.6 | 12.0 | 17.0 | 39.0 | 135.9 |
| Miles Driving | 160.0 | 204.4 | 258.9 | | | | 92.4 | 32.7 | 109.0 | 262.0 | 604.0 | 1723.4 |
| Hours Walking | 23.8 | 46.8 | 180.5 | 29.5 | 3.0 | 2.0 | 52.0 | 34.0 | 16.0 | 34.0 | 52.0 | 473.6 |
| Miles Walking | 17.7 | 38.9 | 133.5 | 20.6 | 2.5 | 3.5 | 36.8 | 28.7 | 13.0 | 17.0 | 22.0 | 334.2 |
| Hours Boating | | | 0.5 | 3.2 | | | 4.5 | | | | | 8.2 |
| Miles Boating | | | 0.4 | 2.1 | | | 5.0 | | | | | 7.5 |
| Stationary Hours | | 0.5 | 9.4 | 2.0 | | 1.7 | | 7.9 | | 4.5 | | 26.0 |
| Feeder Hours | 7.3 | 7.5 | 3.5 | | 2.5 | | 3.0 | 3.0 | | 4.0 | | 30.8 |
| Hours Owling | 2.5 | 0.3 | 2.0 | | | | 1.5 | 0.4 | | 5.0 | 7.0 | 18.7 |
| Miles Owling | | 0.1 | 1.3 | | | | 1.0 | 0.3 | | 20.0 | 1.0 | 23.7 |
| Hours Other | | | | | | | | | | 2.3 | | 2.3 |
| Miles Other | | | | | | | | | | 2.0 | | 2.0 |

There was one species that was a write-in to the distributed checklist for this count, a single Tundra Swan found in Dorchester County. Additional species with only a single individual found statewide included Ruffed Grouse (Allegany), Eastern Whip-poor-will (Allegany), American Coot (Prince George's), American Avocet (Dorchester), Baird's Sandpiper (Allegany), Western Sandpiper (Dorchester), Little Blue Heron (Prince George's), Barn Owl (Calvert), Yellow-bellied Sapsucker (Caroline), Olive-sided Flycatcher (Prince George's), Yellow-bellied Flycatcher (Prince George's), Blue-headed Vireo (Allegany), Prothonotary Warbler (Calvert), and Grasshopper Sparrow (Calvert) (Table 3).

The following highlights were noted by compilers in their inputs. The 60 Red-breasted Nuthatches observed across the state (with only two misses) reflects this was a significant irruption year. Calvert observed that the total absence of swallows was unusual, suggesting that perhaps it was too "sultry" for them, however the presence of swallows seemed to be low for all counts west of the

Chesapeake Bay. In Dorchester, an American Avocet was a first record for their count, the 8 American White Pelicans were of note, the 365 Brown Pelicans were a new high (222 last year), and the 2 Orchard Orioles were a very late record. Despite excellent coverage, significant Dorchester misses were Eastern Meadowlark and Song Sparrow. In Howard, the 4 American Woodcocks were only the second record in 23 counts, and new high counts were Wild Turkey 14, Green Heron 37, Bald Eagle 15, and Eastern Screech-Owl 9.

There were 36 species that were observed in only one county. Fifteen of these were observed in Dorchester County, with the rest being 7 in Allegany, 5 in Calvert, 4 in Prince George's, 2 in Caroline, and 1 each in Harford, Howard, and Kent County. Twenty-two species were reported for all 11 counts.

Thank you to all participants and especially the compilers. I urge more participants to enjoy the experience and join in one of the 2017 counts that will be held on 16 or 17 September. I hope additional counties will have volunteers who choose to organize a count in the future and reverse a trend in decreasing interest and participation in organized seasonal counts.

Table 3. 2016 Maryland Fall Count: observed species.

| Species | AL | CA | HO | BA | HA | CE | PG | CT | KE | CN | DO | Total |
|---------------------------|----|-----|------|----|----|----|-----|----|-----|-----|------|-------|
| Canada Goose | 35 | 589 | 1198 | 61 | 3 | | 581 | 68 | 86 | 171 | 1247 | 4039 |
| Mute Swan | | | | | 2 | | | | | | | 2 |
| Tundra Swan | | | | | | | | | | | 1 | 1 |
| Wood Duck | 8 | | 27 | 28 | 1 | | 45 | 71 | 19 | 20 | 33 | 252 |
| American Wigeon | | | | | | | | | 16 | | | 16 |
| American Black Duck | | 3 | | | | | 2 | 4 | 6 | 4 | 16 | 35 |
| Mallard | 24 | 203 | 154 | 88 | | | 229 | 96 | 873 | 31 | 562 | 2260 |
| Blue-winged Teal | | 1 | | | | | 4 | 2 | 9 | | | 12 |
| Northern Pintail | | | | | | | | | 3 | 5 | 2 | 10 |
| Green-winged Teal | 3 | | | | | | 12 | | 67 | | 111 | 193 |
| Ruddy Duck | | | | | | | | | 1 | | 8 | 9 |
| Northern Bobwhite | | | | | | | | | | 6 | 3 | 9 |
| Ruffed Grouse | 1 | | | | | | | | | | | 1 |
| Wild Turkey | 13 | 2 | 14 | | | | 1 | 1 | | 46 | 6 | 83 |
| Pied-billed Grebe | | 1 | 1 | 1 | | | 8 | | | | | 11 |
| Rock Pigeon | 37 | 127 | 221 | 39 | 1 | | 83 | 3 | 12 | 72 | 13 | 608 |
| Mourning Dove | 64 | 240 | 742 | 44 | 38 | | 299 | 69 | 53 | 199 | 136 | 1884 |
| Yellow-billed Cuckoo | | 4 | 4 | 1 | | | 3 | 7 | | | 4 | 23 |
| Black-billed Cuckoo | | | 1 | | | | | 1 | | | 1 | 3 |
| Common Nighthawk | | | 6 | 1 | | | | | 1 | | | 8 |
| Eastern Whip-poor-will | 1 | | | | | | | | | | | 1 |
| Chimney Swift | 36 | 43 | 182 | 17 | | 4 | 246 | 11 | | 15 | 30 | 584 |
| Ruby-throated Hummingbird | 8 | 34 | 33 | 1 | 3 | 2 | 8 | 16 | 8 | 24 | 8 | 145 |
| Clapper Rail | | | | | | | | | | | 6 | 6 |

| Species | AL | CA | HO | BA | HA | CE | PG | CT | KE | CN | DO | Total | |
|----------------------------|----|----|-----|-----|----|----|-----|------|-----|-----|-----|-------|------|
| Virginia Rail | | | | | | | | | 2 | | 41 | 43 | |
| Sora | | | | | | | 4 | | | | 1 | 5 | |
| Common Gallinule | | | | | | | | | | | 2 | 2 | |
| American Coot | | | | | | | 1 | | | | | 1 | |
| American Avocet | | | | | | | | | | | 1 | 1 | |
| Black-bellied Plover | | | | | | | 1 | | | | 2 | 3 | |
| Semipalmated Plover | | | | 11 | | | 3 | 5 | | | 34 | 53 | |
| Killdeer | 5 | 59 | 44 | 45 | | | 32 | 21 | 16 | 4 | 128 | 354 | |
| Stilt Sandpiper | | | | 4 | | | | | | | 1 | 5 | |
| Sanderling | | | | | | | | 13 | | | 5 | 18 | |
| Dunlin | | | | | | | | | | | 42 | 42 | |
| Baird's Sandpiper | 1 | | | | | | | | | | | 1 | |
| Least Sandpiper | 18 | 7 | 25 | 27 | | | 24 | 1 | 14 | | 85 | 201 | |
| Pectoral Sandpiper | 6 | 1 | | 18 | | | 17 | | | | 20 | 62 | |
| Semipalmated Sandpiper | | | 4 | 14 | | | 5 | 1 | | | 53 | 77 | |
| Western Sandpiper | | | | | | | | | | | 1 | 1 | |
| Short-billed Dowitcher | | | | | | | | | | | 2 | 2 | |
| Wilson's Snipe | | | | | | | | | | 4 | 2 | 6 | |
| American Woodcock | | | 4 | | | | | | | | 1 | 5 | |
| Spotted Sandpiper | | | 1 | 3 | | | 1 | 1 | | 6 | 3 | 15 | |
| Solitary Sandpiper | 1 | | 2 | 3 | | | 2 | | 1 | | | 9 | |
| Greater Yellowlegs | 1 | 1 | | 7 | | | 2 | | 14 | | 25 | 50 | |
| Lesser Yellowlegs | 1 | 1 | 1 | 36 | | | 28 | | 4 | | 20 | 91 | |
| unidentified shorebird | | | | | | | | | | | 100 | 100 | |
| Laughing Gull | | | | 157 | | | 2 | 576 | 791 | 284 | 842 | 1192 | 3844 |
| Ring-billed Gull | | | 9 | 9 | | | 112 | 179 | 36 | 45 | 246 | 636 | |
| Herring Gull | | | | 3 | | | 1 | 186 | 13 | | 381 | 584 | |
| Great Black-backed Gull | | | | 19 | | | 1 | 1464 | 15 | | 67 | 1566 | |
| unidentified gull | | | | 200 | | | | 420 | | 73 | | 693 | |
| Caspian Tern | | | | 2 | | 1 | 10 | 2 | 17 | | 45 | 77 | |
| Black Tern | | | | | | | 2 | | | | | 2 | |
| Common Tern | | | | | | | | 7 | 39 | 9 | 18 | 73 | |
| Forster's Tern | | | | | | | 18 | 351 | 322 | 3 | 331 | 1025 | |
| Royal Tern | | | | | | | 9 | 60 | 30 | | 265 | 364 | |
| Double-crested Cormorant | | 5 | 18 | 50 | | 15 | 48 | 649 | 153 | 14 | 233 | 1185 | |
| American White Pelican | | | | | | | | | | | 8 | 8 | |
| Brown Pelican | | | | | | | | 6 | | | 365 | 371 | |
| Great Blue Heron | 2 | 10 | 35 | 16 | 1 | | 29 | 16 | 26 | 18 | 67 | 220 | |
| Great Egret | | 2 | 9 | | | | 44 | 4 | 3 | 2 | 108 | 172 | |
| Snowy Egret | | | | | | | 5 | 8 | 2 | | 104 | 119 | |
| Little Blue Heron | | | | | | | 1 | | | | | 1 | |
| Tricolored Heron | | | | | | | | | | | 4 | 4 | |
| Cattle Egret | | | | | | | | | | | 40 | 40 | |
| Green Heron | 1 | 11 | 37 | 4 | 1 | | 11 | 1 | 2 | 1 | 4 | 73 | |
| Black-crowned Night-Heron | | | 1 | 1 | | | | | 1 | | 1 | 4 | |
| Yellow-crowned Night-Heron | | | | 1 | | | 1 | | | | | 2 | |
| Glossy Ibis | | | | | | | | | | | 5 | 5 | |
| Black Vulture | 3 | 31 | 106 | 13 | | 3 | 49 | 37 | 16 | 88 | 32 | 378 | |

| Species | AL | CA | HO | BA | HA | CE | PG | CT | KE | CN | DO | Total |
|-------------------------------|-----|-----|-----|----|----|----|-----|-----|-----|-----|-----|-------|
| Turkey Vulture | 24 | 158 | 205 | 29 | 5 | 4 | 76 | 80 | 149 | 240 | 383 | 1353 |
| Osprey | | | 12 | 4 | | 1 | 6 | 26 | 6 | 9 | 29 | 93 |
| Bald Eagle | 3 | 5 | 15 | 7 | 2 | 1 | 8 | 42 | 69 | 20 | 149 | 321 |
| Northern Harrier | 1 | 1 | 1 | | | | 1 | 1 | | 2 | 12 | 19 |
| Sharp-shinned Hawk | 4 | 3 | 6 | | | 2 | 1 | 1 | 3 | 1 | 1 | 22 |
| Cooper's Hawk | 1 | 4 | 15 | 4 | | | 5 | 2 | 1 | 3 | 4 | 39 |
| unidentified <i>Accipiter</i> | | | 2 | | | | | | | | | 2 |
| Red-shouldered Hawk | 2 | 9 | 59 | 6 | | | 16 | 6 | 1 | 3 | 5 | 107 |
| Broad-winged Hawk | 3 | | 3 | | | | 1 | | | | 8 | 15 |
| Red-tailed Hawk | | 15 | 15 | 2 | | | 13 | 3 | 6 | 7 | 20 | 81 |
| unidentified <i>Buteo</i> | 2 | | | | | | | | | | | 2 |
| Barn Owl | | | | | | | | 1 | | | | 1 |
| Eastern Screech-Owl | 1 | 3 | 9 | | 1 | 3 | 3 | 2 | 2 | 5 | 22 | 51 |
| Great Horned Owl | | 1 | 1 | | | | | | 3 | 4 | 24 | 33 |
| Barred Owl | 1 | 2 | 11 | | 1 | | | 4 | 5 | 7 | 2 | 33 |
| Belted Kingfisher | | 26 | 35 | 12 | 1 | 1 | 8 | 7 | 5 | 9 | 14 | 118 |
| Red-headed Woodpecker | | 4 | | | | 1 | 6 | 1 | | | 20 | 32 |
| Red-bellied Woodpecker | 22 | 92 | 208 | 24 | 4 | 2 | 49 | 50 | 49 | 39 | 22 | 561 |
| Yellow-bellied Sapsucker | | | | | | | | | | 1 | | 1 |
| Downy Woodpecker | 12 | 57 | 130 | 19 | 3 | 3 | 35 | 57 | 27 | 36 | 50 | 429 |
| Hairy Woodpecker | 5 | 10 | 26 | 3 | 1 | 3 | 10 | 12 | 9 | 7 | 7 | 93 |
| Northern Flicker | 7 | 55 | 90 | 22 | 1 | 4 | 20 | 13 | 23 | 8 | 35 | 278 |
| Pileated Woodpecker | 17 | 18 | 54 | 6 | 3 | 1 | 14 | 17 | 11 | 22 | 22 | 185 |
| American Kestrel | 2 | 8 | 14 | | | | 17 | 2 | 8 | 9 | 29 | 89 |
| Merlin | | | 2 | | | | | | | | 6 | 8 |
| Peregrine Falcon | | 1 | | | | | | | | | 1 | 2 |
| Olive-sided Flycatcher | | | | | | | 1 | | | | | 1 |
| Eastern Wood-Pewee | 6 | 34 | 59 | 10 | 2 | 2 | 30 | 45 | 28 | 14 | 21 | 251 |
| Yellow-bellied Flycatcher | | | | | | | | 1 | | | | 1 |
| Acadian Flycatcher | | | 3 | | | | 7 | 25 | 1 | 1 | 1 | 38 |
| Willow Flycatcher | | 1 | 1 | | | | | | | | | 2 |
| Traill's Flycatcher | 2 | | 3 | | | | | | 3 | | | 8 |
| Least Flycatcher | | 1 | 2 | 1 | | 1 | | 1 | | | 1 | 7 |
| unidentified <i>Empidonax</i> | | 13 | 9 | | 1 | 1 | | 3 | | | 3 | 30 |
| Eastern Phoebe | 22 | 32 | 39 | 17 | | | 26 | 18 | 3 | 9 | 9 | 175 |
| Great Crested Flycatcher | 1 | 2 | 3 | 2 | | 1 | 2 | 7 | 5 | 1 | 14 | 38 |
| Eastern Kingbird | | | 2 | | 1 | | | 1 | | | 6 | 10 |
| White-eyed Vireo | | 10 | 24 | 1 | | 1 | 18 | 47 | 13 | 8 | 28 | 150 |
| Yellow-throated Vireo | 1 | 4 | 4 | 2 | | | | 8 | 2 | | | 21 |
| Blue-headed Vireo | 1 | | | | | | | | | | | 1 |
| Philadelphia Vireo | 3 | 2 | 2 | 2 | | | | 2 | | | | 11 |
| Warbling Vireo | 1 | | 2 | 2 | | | | 1 | | | | 6 |
| Red-eyed Vireo | 15 | 26 | 46 | 10 | | 5 | 12 | 75 | 22 | 5 | 19 | 235 |
| Blue Jay | 53 | 247 | 609 | 80 | 4 | 21 | 105 | 89 | 68 | 126 | 139 | 1541 |
| American Crow | 104 | 373 | 480 | 52 | 7 | 5 | 99 | 239 | 29 | 133 | 263 | 1784 |
| Fish Crow | | 12 | 44 | 7 | | 45 | 77 | 39 | | 82 | 7 | 313 |
| unidentified crow | | | 311 | 18 | | | 86 | 4 | | 30 | 10 | 459 |
| Common Raven | 10 | | 4 | | | | 1 | | | | | 15 |

| Species | AL | CA | HO | BA | HA | CE | PG | CT | KE | CN | DO | Total |
|-------------------------|-----|------|------|-----|----|----|-----|-----|-----|------|------|-------|
| Horned Lark | | 40 | 10 | | | | | | 25 | 18 | 18 | 111 |
| Tree Swallow | | | 1 | | | | | | 107 | 569 | 779 | 1456 |
| N. Rough-winged Swallow | 4 | | | | | | | | 7 | 4 | 12 | 27 |
| Barn Swallow | | 2 | | | | | | | | | 11 | 13 |
| Carolina Chickadee | | 151 | 269 | 30 | 7 | 5 | 66 | 144 | 74 | 126 | 148 | 1020 |
| Black-capped Chickadee | 35 | | | | | | | | | | | 35 |
| Tufted Titmouse | 31 | 108 | 199 | 18 | 7 | 4 | 55 | 158 | 46 | 98 | 123 | 847 |
| Red-breasted Nuthatch | 2 | 7 | 7 | | 2 | | 5 | 1 | 10 | 5 | 21 | 60 |
| White-breasted Nuthatch | 25 | 59 | 125 | 14 | 4 | 2 | 28 | 44 | 8 | 22 | 1 | 332 |
| Brown-headed Nuthatch | | | | | | | | 5 | 4 | | 77 | 86 |
| Brown Creeper | | | 1 | | | | | | | 2 | | 3 |
| House Wren | | 15 | 31 | 9 | 1 | 4 | 14 | 10 | 14 | 1 | 27 | 126 |
| Winter Wren | | | 1 | 1 | | | | | | | | 2 |
| Marsh Wren | | | | | | | | | | | 8 | 8 |
| Carolina Wren | 23 | 96 | 291 | 35 | 2 | 7 | 84 | 109 | 73 | 77 | 112 | 909 |
| Blue-gray Gnatcatcher | | 7 | 17 | | | 1 | 6 | 7 | 4 | 2 | 27 | 71 |
| Golden-crowned Kinglet | 6 | | | | | | | | | | | 6 |
| Ruby-crowned Kinglet | | 1 | | | | | | 1 | | | | 2 |
| Eastern Bluebird | 22 | 177 | 211 | 21 | 6 | | 48 | 52 | 40 | 45 | 64 | 686 |
| Veery | 1 | 1 | 5 | 1 | | | | 14 | 8 | | | 30 |
| Gray-cheeked Thrush | | 1 | 1 | | | | | | | | | 2 |
| Swainson's Thrush | 5 | | 6 | | | | | 1 | 4 | | | 16 |
| Hermit Thrush | | 1 | | | | | | | | 3 | | 4 |
| Wood Thrush | 2 | 13 | 28 | 4 | | | 4 | 9 | 13 | 2 | 3 | 78 |
| American Robin | 24 | 430 | 479 | 74 | | | 170 | 31 | 17 | 140 | 402 | 1767 |
| Gray Catbird | 16 | 273 | 343 | 45 | 2 | 6 | 64 | 25 | 33 | 13 | 34 | 854 |
| Brown Thrasher | 5 | 9 | 6 | 2 | | 1 | 10 | 12 | 24 | 2 | 16 | 87 |
| Northern Mockingbird | 4 | 59 | 82 | 5 | 3 | | 37 | 17 | 38 | 60 | 52 | 357 |
| European Starling | 77 | 1319 | 2572 | 119 | | | 594 | 37 | 235 | 1211 | 1099 | 7263 |
| Cedar Waxwing | 305 | 83 | 93 | | 2 | 20 | 21 | | 5 | 21 | 14 | 564 |
| House Sparrow | 41 | 219 | 286 | 36 | 1 | | 183 | 32 | 75 | 168 | 34 | 1075 |
| House Finch | 12 | 122 | 153 | 7 | 12 | 1 | 58 | 26 | 23 | 40 | 4 | 458 |
| Purple Finch | 3 | | | | | | | | | | | 3 |
| American Goldfinch | 93 | 178 | 490 | 69 | 18 | 3 | 78 | 54 | 58 | 53 | 49 | 1143 |
| Ovenbird | | 2 | 3 | 1 | 1 | 1 | | 1 | | | 2 | 11 |
| Worm-eating Warbler | | | 1 | | | | | 1 | | | | 2 |
| Northern Waterthrush | | | 6 | | | | | 5 | 1 | | 2 | 14 |
| Blue-winged Warbler | | | 1 | | | | 3 | | | | | 4 |
| Black-and-white Warbler | 1 | 9 | 29 | 4 | 2 | 1 | 12 | 17 | 10 | 7 | 13 | 105 |
| Prothonotary Warbler | | | | | | | | 1 | | | | 1 |
| Tennessee Warbler | 4 | 2 | 2 | 1 | | 3 | | | | | | 12 |
| Nashville Warbler | 1 | | 1 | | | | | | | | | 2 |
| Common Yellowthroat | 7 | 54 | 119 | 22 | 5 | 9 | 31 | 40 | 43 | 14 | 49 | 393 |
| Hooded Warbler | | | 1 | | | | | 8 | | | | 9 |
| American Redstart | 2 | 21 | 73 | 6 | 3 | 1 | 15 | 27 | 36 | 23 | 48 | 255 |
| Cape May Warbler | 6 | | | | | 2 | | 1 | 1 | | 2 | 12 |
| Northern Parula | | 5 | 18 | 2 | 1 | 2 | 6 | 9 | 9 | 4 | 6 | 62 |
| Magnolia Warbler | 10 | 6 | 35 | 6 | 1 | | 6 | 8 | 9 | 7 | 3 | 91 |

| Species | AL | CA | HO | BA | HA | CE | PG | CT | KE | CN | DO | Total |
|--------------------------------|----|-----|-----|----|-----|----|------|-----|-----|------|------|-------|
| Bay-breasted Warbler | 2 | 1 | | | | | | | | | | 3 |
| Blackburnian Warbler | 3 | 2 | 2 | | | | | | | | | 7 |
| Yellow Warbler | | | 3 | | | | | 2 | | 3 | 5 | 13 |
| Chestnut-sided Warbler | | 2 | 10 | 4 | | | 4 | 4 | 4 | | | 28 |
| Blackpoll Warbler | 4 | | | | | | 2 | | | | 1 | 7 |
| Black-throated Blue Warbler | 5 | 6 | 9 | 1 | | 1 | 1 | 7 | | 2 | 1 | 33 |
| Palm Warbler | | | 10 | 5 | 12 | | 2 | 2 | 3 | 3 | 5 | 42 |
| Pine Warbler | | 4 | 2 | 2 | | | 1 | 15 | 7 | 6 | 86 | 123 |
| Yellow-rumped Warbler (Myrtle) | | | | | | | | | | 7 | | 7 |
| Yellow-throated Warbler | | | | | | | | 3 | | | | 3 |
| Prairie Warbler | 1 | | 2 | | 2 | | | | 2 | | 3 | 10 |
| Black-throated Green Warbler | 22 | 9 | 10 | | | | 1 | 2 | 3 | 1 | 1 | 49 |
| Canada Warbler | 1 | | 4 | | | | | | | | | 5 |
| Wilson's Warbler | | | 2 | 2 | 1 | | | | | | | 5 |
| Yellow-breasted Chat | | 2 | | | | | | 3 | | | | 5 |
| unidentified warbler | | | 10 | 1 | | 3 | 1 | | | | 8 | 23 |
| Eastern Towhee | 13 | 23 | 45 | | | 1 | 8 | 22 | 5 | 4 | 9 | 130 |
| Chipping Sparrow | 38 | 57 | 191 | 21 | 23 | | 48 | 120 | 90 | 68 | 55 | 711 |
| Field Sparrow | 9 | 4 | 10 | | | | 1 | 4 | 7 | 8 | 3 | 46 |
| Savannah Sparrow | | 1 | 3 | | 1 | | | | | | 5 | 10 |
| Grasshopper Sparrow | | | | | | | | 1 | | | | 1 |
| Seaside Sparrow | | | | | | | | | | | 14 | 14 |
| Song Sparrow | 14 | 34 | 57 | 14 | 2 | | 10 | 5 | 1 | 2 | | 139 |
| Swamp Sparrow | | | 1 | 1 | | | | | | | 1 | 3 |
| unidentified sparrow | | | | | | | | | | | 1 | 1 |
| Summer Tanager | | | | | | | 3 | 12 | | 2 | 5 | 22 |
| Scarlet Tanager | 19 | 8 | 8 | 4 | 2 | 1 | 5 | 7 | 14 | 5 | 1 | 74 |
| Northern Cardinal | 41 | 217 | 397 | 58 | 10 | 11 | 143 | 183 | 155 | 132 | 86 | 1433 |
| Rose-breasted Grosbeak | 5 | 4 | 5 | 1 | | | 3 | 3 | | | | 21 |
| Blue Grosbeak | | | 6 | 3 | 8 | | 6 | 15 | 12 | 37 | 61 | 148 |
| Indigo Bunting | 3 | 9 | 91 | | 9 | | 19 | 52 | 22 | 30 | 9 | 244 |
| Bobolink | | 1 | 75 | | | 2 | | 1 | 51 | 1 | 717 | 848 |
| Red-winged Blackbird | | 370 | 109 | 24 | | | 22 | 173 | 4 | 117 | 1934 | 2753 |
| Eastern Meadowlark | | | 12 | | | | | | | | | 12 |
| Common Grackle | 30 | | 6 | 45 | 300 | | 3064 | 14 | 26 | 1050 | 291 | 4826 |
| Boat-tailed Grackle | | | | | | | | | | | 8 | 8 |
| Brown-headed Cowbird | | 20 | 138 | | 3 | | 80 | 5 | | 125 | 942 | 1313 |
| unidentified blackbird | | | | | | | 1 | | | | | 1 |
| Orchard Oriole | | | | | | | | | | | 2 | 2 |
| Baltimore Oriole | | 2 | 2 | | | | 2 | 1 | | | 1 | 8 |

MARYLAND BIRDLIFE

NOTICE TO CONTRIBUTORS

Contributors should prepare manuscripts according to the following instructions.

Title: The title should be brief, concise, and pertinent.

Abstract: An abstract is required for all long articles; suggested for all biologic studies more than two (2) pages in length; but is not needed for notes, distribution reports, or short observations (especially if two pages or shorter in length). The abstract should provide a capsule description of the main thrust, methods and essential findings of the article. It should contain the scientific name of the main subject species.

Text: Manuscripts should be double-spaced, lines numbered, and submitted in MS Word™ by e-mail or on a CD. Please identify respective file name(s) for text, figure titles, and descriptions of graphs or figures. First mention of a biological organism, in the abstract and text should include the full scientific name in italics. Carefully check the spelling of all scientific names. Capitalize the first letter of each word comprising the “official” common name for faunal species. Short articles and general notes (20 pages or less) are preferred. Color “copy ready” illustrations, pictures, or digital images are preferred.

References: References should be given in an author-date format: (Robbins 1987); (Robbins 1987, 1988); (Robbins, in press); (Robbins, in litt.); (Robbins, pers. comm.); (Robbins and Robbins 1987); and (Robbins et al. 1987) for three or more authors. Provide evidence of acceptance for works “in press,” or cite as “unpublished”, “in litt.”, or “pers. comm.” (written permission is suggested as well). Citations shall be listed alphabetically, under LITERATURE CITED, as follows: **Articles:** Robbins, C.S. 1965. New breeding bird survey tested in Maryland this summer. *Maryland Birdlife* 21(2):48-49. Do not abbreviate the titles of journals. **Books:** Ellison, W.G. (Editor). 2010. *2nd Atlas of the Breeding Birds of Maryland and the District of Columbia*. The Johns Hopkins University Press, Baltimore, MD. 494 pp. **Internet:** Holt, D.W., M.D. Larson, N. Smith, D.L. Evans, and D.F. Parmalee. 2015. Snowy Owl (*Bubo scandiacus*), The Birds of North America (P.G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY; Available at: <https://birdsna.org/Species-Account/bna/species/snoow11>. Accessed 6 April 2017.

Tables: Tables, graphs and line drawings should be created electronically in black and white. Color should only be used when absolutely necessary for clarity.

Illustrations: Photographs or high-definition images may be accepted if necessary or desired by the author(s) to support the text. Photographs should be submitted in color. Figure numbers, as cited in the text, and figure legends should be keyed to each respective photograph.

Maryland Birdlife is published twice annually to record and encourage the study of birds in and around Maryland. *Maryland Birdlife* contains original articles, notes, and research papers primarily pertaining to Maryland and the Mid-Atlantic region. Potential topics may include geographic or temporal distribution, ecology, biology, morphology, systematics, behavior, migration, life history, as well as other biological topics. Annual bird counts also will be published. All submissions are subject to editorial review and acceptance. Articles and research papers will be peer-reviewed. Send submissions to the Editor: Eugene J. Scarpulla, 14207 Lakerun Court, Bowie, MD 20720-4861, ejscarp@comcast.net.

Non-Profit Org.
U.S. Postage
PAID
Hagerstown, MD
Permit No. 184

Maryland Ornithological Society, Inc.
P.O. Box 105
Monrovia, MD 21770-0105



Maryland Birdlife

Published Semiannually by the Maryland Ornithological Society, Inc.

Editor: Eugene J. Scarpulla, 14207 Lakerun Court, Bowie, MD 20720-4861
ejscarp@comcast.net
Associate Editor: Mark S. Johnson, 3204 Bryson Court, Baldwin, MD 21013
MarkSJohnson2@gmail.com

CONTENTS – SPRING 2017

Editor's Note

| | |
|---|----|
| <i>Eugene J. Scarpulla</i> | 1 |
| Long-eared Owl (<i>Asio otus</i>) Annual Frequency, Seasonal Chronologies, and Status on the Delmarva Peninsula | |
| <i>Jan G. Reese and Robbie Weterings</i> | 2 |
| Novel Winter Bathing Behavior by a Merlin (<i>Falco columbarius</i>) | |
| <i>David R. Smith</i> | 20 |
| First Maryland Record of Pacific-slope/Cordilleran ("Western") Flycatcher, <i>Empidonax difficilis/occidentalis</i> | |
| <i>Kathleen B. Calvert</i> | 23 |
| Sunbathing by Great Crested Flycatchers, <i>Myiarchus crinitus</i> | |
| <i>Barbara K. Johnson</i> | 31 |
| Cedar Waxwings (<i>Bombycilla cedrorum</i>) Observed Feeding on Flying Insects | |
| <i>Jan G. Reese</i> | 39 |
| 2016 Maryland Mid-winter Bird Count: Corrigendum | |
| <i>Mary Anne Fluke</i> | 47 |
| 2016 Maryland May Count | |
| <i>Marilyn E. Veek</i> | 48 |
| 2016 Maryland Fall Count | |
| <i>Charles R. Stirrat</i> | 62 |